

Preceding crops and residue management practices on performance of mustard under zero-till semi-arid condition

LP Amgain¹ and AR Sharma²

¹Institute of Agriculture and Animal Sciences, Rampur, Chitwan, Nepal

²Directorate of Weed Science Research, Jabalpur, India

Abstract

A field experiment was conducted in 2010/11 and 2011/12 at New Delhi to study the performance of zero-till rainfed mustard. The influence of preceding rainy season crops, viz. pearl millet, cluster bean and green gram; and residue management, viz. no residue, crop residue and *Leucaena* twigs mulching on this mustard crop was observed. Pooled analysis of mustard seed yield was significantly higher (+51%) in 2010/11 (1.80 t/ha) than 2011/12 (1.19 t/ha) due to favourable weather condition in the first year. Further, *Leucaena* twigs mulching resulted higher seed yield of 2.17 t/ha in 2010/11 than 1.94 t/ha in 2011/12. Green gram as preceding crop resulted significantly higher mustard yield in 2010/11, while it was higher after cluster bean in 2011/12. Interaction between crop residue and preceding rainy season crops on growth parameters exerted significant variations, while yield attributes showed the mixed response. Cluster bean with *Leucaena* twigs led to higher yield of mustard, followed by green gram with crop residue. Economic analysis exhibited the highest net return of mustard grown after cluster bean with *Leucaena* twigs mulching. From the findings it is observed that growth indices, yield, and yield attributes and economics of mustard were influenced significantly by preceding crops and crop residue application. The effect of *Leucaena* twigs was found better in 2010/11, while both crop residues and *Leucaena* twigs mulching were equally effective in 2011/12. Cluster bean as preceding crop to mustard resulted higher yield and net returns, followed by green gram and cluster bean with crop residues. It was suggested that mustard after cluster bean with *Leucaena* twigs was a high yielding and profitable cropping system under zero-till semi-arid condition. It was concluded that growing mustard after cluster bean with *Leucaena* twigs mulching resulted increased yield and profitable cropping system for rainfed areas in zero-till semi arid condition of New Delhi.

Key words : economics, preceding crops, rainfed mustard, residue management, zero-till

Introduction

Fallowing or cultivation of short duration crops like pearl millet [*Pennisetum glaucum* (L.) R Br Emend Stuntz], cluster bean [*Cyamopsis tetragonoloba* (L.) Taub.], and green gram (*Vigna radiata* L.) during rainy season followed by a long duration drought hardy crop like mustard (*Brassica juncea* L. Czern. & Coss.) during winter season on the conserved soil moisture is commonly followed in semi-arid areas of India and Pakistan (Faroda *et al.*, 2007; Singh *et al.*, 2008). These crops are mostly grown in ploughed land, which not only deteriorates the soil environment, but also increases the cost of production. Conservation tillage (zero-tillage + residue mulching + crop diversification) is a useful practice in rainfed areas to control soil erosion and weed growth as well as to preserve soil moisture and plant nutrients in the soil profile (Narain and Singh, 1997; Sharma *et al.*, 2005a). Inclusion of legumes in the crop rotation further plays a vital role in improving

balanced plant nutrition (Saxena, 2012) and building up of soil fertility through the addition of biologically fixed nitrogen (Ali *et al.*, 2002).

There are several evidences of remarkable increase in the yield of crops in the rainfed cropping system through the maintenance of appropriate vegetative cover under no-till condition (Dhyani *et al.*, 2009). Mulching of crop field by *Leucaena leucocephala* helps in conserving soil moisture for proper growth and development of crops (Sharma *et al.*, 2010). The use of vegetative cover under zero-tillage condition helps to increase root growth by creating favourable soil environment and decreasing weed infestation. This situation is ideal for better plant growth and higher yield of winter crops (Singh *et al.*, 1998). *In-situ* application of the residue of pearl millet, cluster bean and green gram in winter season crops, and the residue of mustard in rainy season crops is practiced because of the easy access to residual material through seasonal harvest. In zero-till condition, the introduction of happy seeder (a machine) makes it easy to sow seeds of any crop, like mustard in standing residue (Jat *et al.*, 2009). This discussion suggests that the adoption of resource conserving technologies, such as zero-tillage and residue management is essential in rainfed condition to improve productivity, resource-use efficiency and sustainability of low input agriculture. Therefore, the aim in this research is to understand the effects of preceding crops and residue management in mustard under zero-till semi-arid condition.

Materials and methods

A field experiment was conducted in 2010/11 and 2011/12 at the Indian Agricultural Research Institute, New Delhi (28° 40'N, 77°12'E) at an altitude of 228 m above the mean sea level in sandy-loam soil to study the effects of preceding rainy season crops and residue management on growth, productivity, nutrient uptake, and economics of mustard-based cropping system. The soil of the research site contains 147.2 kg/ha alkaline KMnO₄-oxidizable N, 17.0 kg/ha NaHCO₃-extractable P, 225.1 kg/ha 1N NH₄OAc-exchangeable K, 0.40% organic C with 7.5 pH (1:2.5 soil and water ratio). The area received 954 mm rainfall in 2010/11. In 2011/12, the occurrence of rainfall in the same area was 662 mm, which was 30.6% less than that of 2010/11 and 10.4% less than the average of the previous 10 years (739 mm). The number of rainy days and the volume of rainfall both were higher during the mustard growing season (October to March) in 2010/11 than in 2011/12. In 2010/11, the number of rainy days during the mustard growing season was 10 and the volume of rainfall was 85 mm, but in 2011/12 the number of rainy days was two and the volume of rainfall was 14 mm. Mustard crop sown on 3rd October 2011 did not germinate for 25 days and to ensure the germination, irrigation measuring 20 mm was given on crop rows. Three cropping systems with combinations of three crops each (pearl millet, cluster bean and green gram) were grown in sequence in rainy season before mustard, exclusively under zero-till rainfed condition following other recommended package of practices with three treatments of surface cover management, viz. control (no residue), crop residues @ 5 t/ha and *Leucaena* twigs @ 10 t/ha.

The experiment was laid out in Randomized Block Design with four replications. Mustard cv. 'Pusa Vijaya' was sown on 18 October 2010, and 3 October 2011 at 40 cm row spacing by using happy seeder. The crop was fertilized with 60.40.20 kg NPK per hectare. The full dose of P and K along with half N was applied through DAP, MOP and urea in the basal application. Diammonium phosphate was mixed with seeds of mustard and placed together in seed box of happy seeder for its proper distribution. However, muriate of potash and urea were broadcasted in the field. Crops were matured from the second to third week of March in both years.

Growth indices, like crop growth rate (CGR), relative growth rate (RGR), and net assimilation rate (NAR) were calculated from 30 DAS to harvest. The yield attributes like plant population per square metre was counted at maturity from the fixed one metre row inserted with pegs from the beginning, while primary branches per plant and siliquae per plant were counted from randomly selected five plants. Number of seeds per siliqua and 1000 seed weight were taken from randomly selected 20 siliquae. The seed and stover yields, and harvest index were recorded from a plot of 10 m² area. The recorded seed yield was adjusted at 10% for moisture. Pooled analysis of seed yield was carried out for evaluation of year effect. Economic analysis was undertaken to find out the cost of cultivation, gross and net returns, and net returns on investment. The biometric data on ancillary and yield parameters were analyzed by using standard statistical techniques. The regression analysis of major yield attributes and seed yield was also done.

Results and discussion

Soil moisture availability and growth indices of mustard


The calculated data showed that the values of CGR, RGR, and NAR between 30-60 DAS and 60-90 DAS were higher under the *Leucaena* twigs than the crop residue after cluster bean and green gram as preceding crops (Table 1 and Table 2). A comparatively higher CGR was recorded between 0 to 30 DAS in 2010/11. This might be due to 22 and 10 mm of rain received within two to three weeks of sowing. The soil moisture condition was different to produce different result in 2011/12. The initial soil moisture content was very low due to high evaporation rate coinciding with high ambient temperature during October 2011. Due to this reason, shallow irrigation in crop rows could not improve the growth of the crop. The moisture availability was directly influencing the CGR, RGR and NAR at different stages of mustard in 2010/11 and 2011/12 (Figure 1).

Table 1. Effect of crop residue and *Leucaena* twigs on crop growth indices of mustard after rainy season crops in 2010/11

Treatment	CGR (g/day/m ²)				RGR (g/g/day)			NAR (g/day/m ²)	
	0-30 DAS	30-60 DAS	60-90 DAS	90-120 DAS	30-60 DAS	60- 90 DAS	90- 120 DAS	30-60 DAS	60-90 DAS
PM- NR	1.68	2.11	3.05	2.76	0.060	0.065	0.064	0.526	0.318
PM - CR	2.24	4.14	4.61	3.03	0.070	0.071	0.065	0.523	0.332
PM - LT	5.14	4.38	4.00	3.99	0.071	0.069	0.069	0.440	0.360
CB - NR	2.75	2.81	6.72	1.14	0.064	0.077	0.051	1.262	0.200
CB - CR	4.02	3.75	8.45	1.99	0.068	0.080	0.059	0.930	0.170
CB - LT	5.67	4.58	11.04	2.65	0.071	0.084	0.063	0.392	0.150
GG - NR	3.31	1.59	5.06	3.30	0.056	0.073	0.067	1.251	0.222
GG - CR	3.23	3.92	7.51	7.08	0.069	0.078	0.078	0.753	0.182
GG - LT	5.40	2.03	8.08	5.86	0.059	0.079	0.075	0.625	0.180

Table 2. Effect of crop residue and *Leucaena* twigs on crop growth indices of mustard after rainy season crops in 2011/12

Treatment	CGR (g/day/m ²)				RGR (g/g/day)			NAR (g/day/m ²)	
	0-30	30-60	60- 90	90-120	30-60	60-	90-	30- 60	60-90
	DAS	DAS	DAS	DAS	DAS	90	120	DAS	DAS
PM- NR	3.81	1.79	4.23	2.26	0.058	0.070	0.061	0.619	0.601
PM - CR	5.03	3.80	3.19	7.04	0.069	0.066	0.077	0.307	0.923
PM - LT	4.34	2.95	4.54	3.51	0.065	0.071	0.067	0.561	0.511
CB - NR	2.75	2.68	3.15	5.37	0.064	0.066	0.074	0.403	1.460
CB - CR	3.23	5.46	3.32	4.90	0.074	0.067	0.072	0.269	0.828
CB - LT	3.76	6.54	8.78	3.34	0.076	0.081	0.067	0.238	0.486
GG - NR	2.01	2.85	3.06	3.44	0.064	0.065	0.067	0.408	1.317
GG - CR	3.05	5.11	4.93	4.25	0.073	0.072	0.070	0.281	1.030
GG - LT	2.47	4.60	3.61	3.65	0.071	0.068	0.068	0.305	1.753






Figure 1. Soil moisture profile (w/w %) in mustard field as influenced by residue retention practices in 2010/11 and 2011/12

Mustard is a drought hardy crop. This crop can grow and give some yield in rainfed area by utilizing the conserved soil moisture. *Leucaena* twigs and crop residue mulching plots recorded more soil moisture conservation than no residue and helped to augment crop growth. Mustard being grown as rainfed, followed no definite trend in growth indices due to variable availability of soil moisture through rainfall. Forty nine mm of rainfall received in February 2011 and 14 mm of rainfall received in January 2012 helped to augment the CGR in later stages of the crop as it was rejuvenated after getting soil moisture. Crop residue of most crops and *Leucaena* twigs in cluster bean provided relatively higher CGR, RGR and NAR than without residue and after green gram and cluster bean as preceding crops. This indicated that soil environment could be made favourable through the application of crop residue and growing legumes as preceding crops in mustard based system. Several researchers (Rathore *et al.*, 1998; Singh *et al.*, 2003; Rathore *et al.*, 2008) reported higher crop growth rate and RGR of mustard when it is followed by legumes and mulched with crop residue.

Yield and yield attributes of mustard

Yield of mustard, and their correlation and regression with major yield attributes as affected by preceding rainy season crops and residue management practices are presented in Figure 2 and 3. Except plant population per square metre at maturity in 2010/11, the effect of preceding crops on most of the yield attributes, seeds per siliqua in 2011/12, and 1000 seed weight in both years showed significant variation. Legumes improve soil fertility and conserve soil moisture, which led to more yield than pearl millet as preceding crop. All major yield attributes, such as plant population per square metre at maturity, primary branches per plant, siliquae per plant and 1000 seed weight were found to be significantly higher in the plots where residue management was done properly. Cluster bean as previous crop with *Leucaena* twigs as mulch, and green gram and pearl millet as previous crops with crop residue as mulch resulted significantly high plant population per square metre at maturity and 1000 seed weight in 2011/12, and siliquae per plant in both years. Higher yield attributes were observed from the crop mulched with *Leucaena* twigs and crop





Figure 2. Yield performance of mustard as influenced by residue management and preceding rainy season crops






Figure 3. Regression and correlation of mustard yield (y) with major yield attributes (x)

residue. Application of *Leucaena* twigs over two years in the fixed plots increased fertility status and moisture holding capacity of the soil resulting high plant population per square metre (20.3) and 1000 seed weight (4.92 g). The favourable improvements in yield attributes could be attributed to the influence of previous legume crops and organic mulches as a result of greater nutrient uptake, efficient partitioning of metabolites and adequate accumulation of translocation of photosynthates. Adequate supply of moisture enhances the growth and dry matter production of crops (Tetarwal and Rana, 2006, Parihar *et al.*, 2010). Singh *et al.* (2003) and Singh *et al.* (2008) found significant increase in yield attributes of mustard grown after cluster bean and green gram and mulched with crop residue after legume crops.

Regression analysis between yields and major yield attributes of mustard revealed significantly positive correlation between mustard yield and number of siliquae per plant, but non-significant between yield and primary branches per plant (Figure 3).

Pooled analysis of mustard seed yield as affected by years, preceding crops and residue management is presented in Table 3. This Table shows that all production factors affect mustard yield singly as well as in combination. There was 51% higher yield (1.80 ton per hectare) in 2010/11 than in 2011/12 (1.20 ton per hectare) due to favorable weather condition.

Table 3. Pooled analysis on seed yield of mustard ($t\ ha^{-1}$) as affected by year, preceding crops and residue management

Preceding crops	2010/11				2011/12				Overall mean
	NR	CR	LR	Mean	NR	CR	LR	Mean	
Pearl millet	1.34	1.81	2.25	1.80	0.74	1.38	0.99	1.04	1.42
Cluster bean	0.74	2.11	2.29	1.71	0.96	1.26	1.93	1.38	1.55
Green gram	1.49	2.23	1.98	1.90	0.81	1.56	1.13	1.17	1.53
Mean	1.19	2.05	2.17		0.84	1.40	1.35	1.20	
	Year (A)	Preceding crop (B)		Residue (C)	A x B	A x C	B x C	A x B x C	
SEm±	0.037	0.046		0.046	0.065	0.065	0.079	0.112	
CD (P=0.05)	0.075	0.093		0.093	0.131	0.131	0.161	0.227	

There was fair distribution of rainfall throughout the mustard growing season in 2010/11, and the last rainfall (49 mm) received in mid-February was coincided with flowering and fruiting. The use of crop residue controlled evaporation loss and increased soil fertility through its decomposition. As a result, the yield was higher with *Leucaena* twigs than no residue.

Economics of mustard

Economic analysis of mustard revealed that cost of cultivation was relatively higher in 2011/12 than 2010/11 (Table 4). The increase in production cost was due to increase in labour wages from Rs 150/man day in 2010/11 to Rs 200/man day in 2011/12, and costs of other inputs. Another reason of high cost of production was due to the estimation of the value of crop residue in added market price. Although the output price was higher in 2011/12, the net return from the crop was low due to poor yield. The highest returns and net returns on investment were achieved under cluster bean with *Leucaena* twigs. Growing mustard after cluster bean and green gram with crop residue and *Leucaena* twigs helped improving profitability under zero-till semi-arid condition. These findings are in accordance with Saxena *et al.* (1998); Singh *et al.* (2003); and Singh *et al.* (2008).

Table 4. Effect of crop residue and *Leucaena* twigs on economics of mustard after rainy season crops

Treatment	Cost of cultivation (x10 ³ `ha ⁻¹)		Gross returns (x10 ³ `ha ⁻¹)		Net returns (x10 ³ `ha ⁻¹)		Net returns/` invested	
	2010/11	2011/12	2010/11	2011/12	2010/11	2011/12	2010/11	2011/12
Pearl millet - no residue	10.23	14.26	26.12	19.91	15.89	5.65	1.55	0.40
Pearl millet - crop residue	12.68	17.36	36.36	37.75	23.68	20.40	1.87	1.18
Pearl millet - <i>Leucaena</i> twigs	11.73	16.26	44.37	27.05	32.64	10.79	2.78	0.66
Cluster bean - no residue	10.23	14.26	15.04	25.38	4.81	11.12	0.47	0.78
Cluster bean - crop residue	12.68	17.36	41.86	34.49	29.18	17.14	2.30	0.99
Cluster bean - <i>Leucaena</i> twigs	11.73	16.26	45.75	51.56	34.02	35.30	2.90	2.17
Green gram - no residue	10.23	14.26	29.19	21.34	18.96	7.08	1.85	0.50
Green gram - crop residue	12.68	17.36	44.21	42.21	31.53	24.85	2.49	1.43
Green gram - <i>Leucaena</i> twigs	11.73	16.26	39.27	30.60	27.54	14.35	2.35	0.88

Acknowledgements

This work is a part of Ph D study of the first author. ICCR is thanked for providing SAARC scholarship and CIMMYT-India for research grants and field workers of Agronomy Division, IARI, Pusa, New Delhi, India for their precise works.

References

- Ali, M, AN Ganeshmurthy and Ch Srinivasarao. 2002. Role of pulses in soil health and sustainable crop production. *Indian Journal of Pulse Research* 15 (2). 107–117.
- Dhyani, SK, R Newaj and AR Sharma. 2009. Agro-forestry. its relation with agronomy, challenges and opportunities. *Indian Journal of Agronomy* 54 (3). 259–266.
- Faroda, AS, N L Joshi, R Singh and A Saxena. 2007. Resource management for sustainable crop production in arid zone. a review. *Indian Journal of Agronomy* 52 (3). 181–193.
- Gomez, KA and AA Gomez. 1984. *Statistical procedures for agricultural research*. John Willey and Sons, NY.
- Jat, ML, MK Gathala, JK Ladha, YS Saharawat, AS Jat, SK Sharma, V Kumar, and R Gupta. 2009. Evaluation of precision land levelling and double zero-till systems in the rice–wheat rotation. water use, productivity, profitability and soil physical properties. *Soil and Tillage Research* 105. 112–121.

- Parihar, CM, KS Rana and SR Kantwa. 2010. Nutrient management in pearl millet (*Pennisetum glaucum*) – mustard (*Brassica juncea*) cropping system as affected by land configuration under limited irrigation. *Indian Journal of Agronomy* 55 (3). 191-196.
- Rathore, AL, AR Pal and KK Sahu. 1998. Tillage and mulching effects on water use, root growth and yield of rainfed mustard and chickpea grown after lowland rice. *Journal of the Science of Food and Agriculture* 78(2). 149–161.
- Rathore, BS, VS Rana and RK Nanwal. 2008. Effect of plant density and fertility levels on growth and yield of pearl millet (*Pennisetum glaucum*) hybrids under limited irrigation conditions in semi-arid environment. *Indian Journal of Agricultural Sciences* 78 (1). 667–670.
- Saxena, A. 2008. Effect of mustard residues on weed population, soil moisture and cluster bean productivity. *Annals of Arid Zone* 47 (1). 25–32.
- Saxena, A, DV Singh and NL Joshi. 1998. Effects of tillage and cropping systems on soil moisture balance and pearl millet yield. *Journal Agronomy and Crop Science* 178. 251–257.
- Saxena, MC. 2012. Challaneges and opportunities for food legume reeararch and development. Paper presented on the 6th M.S. Swaminathan Award Lecture at IARI, Pusa, New Delhi, January 25, 2012.
- Sharma, AR, R Singh, SK Dhyani and RK Dube. 2010. Moisture conservation and nitrogen recycling through legume mulching in rainfed maize (*Zea mays*)–wheat (*Triticum aestivum*) cropping system. *Nutrient Cycling in Agro-ecosystems* 87(2). 187–197.
- Singh, BP, MC Mundra and SC Gupta. 2003. Productivity, stability and economics of various cropping systems in semi-arid ecosystem. *Crop Research (Hisar)* 25(3). 472-477.
- Singh, R, B Singh and M Patidar. 2008. Effect of preceding crops and nutrient management on productivity of wheat (*Triticum aestivum*) – based cropping system in arid region. *Indian Journal of Agronomy* 53 (4). 267–272.
- Singh, V, SK Sharma, D Ram, RK Siag and BL Verma. 1998. Performance of different crops sequences under various irrigation levels. *Indian Journal of Agronomy* 35. 287–296.
- Tetarwal, JP and KS Rana. 2006. Impact of cropping system, fertility level and moisture conservation practice on productivity, nutrient uptake, water use and profitability of pearl millet (*Pennisetum glaucum*) under rainfed conditions. *Indian Journal of Agronomy* 51(4). 263–266.