



# International Journal of Applied Sciences and Biotechnology

A Rapid Publishing Journal

ISSN 2091-2609



**Available online at:**

<http://www.ijasbt.org>

&

<http://www.nepjol.info/index.php/IJASBT/index>

## **Indexing and Abstracting**

CrossRef, Google Scholar, Global Impact Factor, Genamics, Index Copernicus, Directory of Open Access Journals, WorldCat, Electronic Journals Library (EZB), Universitätsbibliothek Leipzig, Hamburg University, UTS (University of Technology, Sydney): Library, International Society of Universal Research in Sciences (EyeSource), Journal Seeker, WZB, Socolar, BioRes, Indian Science, Jadoun Science, Jour-Informatics, Journal Directory, JournalTOCs, Academic Journals Database, Journal Quality Evaluation Report, PDOAJ, Science Central, Journal Impact Factor, NewJour, Open Science Directory, Directory of Research Journals Indexing, Open Access Library, International Impact Factor Services, SciSeek, Cabell's Directories, Scientific Indexing Services, CiteFactor, UniSA Library, InfoBase Index, Infomine, Getinfo, Open Academic Journals Index, HINARI, etc.

**CODEN (Chemical Abstract Services, USA): IJASKD**

Vol-2(2) June, 2014

IC Value: 4.37



For any type of query and/or feedback don't hesitate to email us at: [editor.ijasbt@gmail.com](mailto:editor.ijasbt@gmail.com)



Research Article

## DETECTION OF TOXIGENIC FUNGI AND MYCOTOXINS IN SOME STORED MEDICINAL PLANT SAMPLES

Ashish Rawat<sup>1</sup>, Surabhi Mahajan<sup>2\*</sup>, Ankur Gupta<sup>2</sup>, Rajneesh Kumar Agnihotri<sup>1</sup>, Nitin Wahi<sup>3</sup> and Rajendra Sharma<sup>1</sup>

<sup>1</sup>Department of Botany, School of Life Sciences, Dr. B.R. Ambedkar University, Agra

<sup>2</sup>Department of Microbiology, School of Life Sciences, Dr. B.R. Ambedkar University, Agra

<sup>3</sup>Department of Biotechnology, GLA University, Mathura

\*Corresponding author email: smahajanmb@gmail.com

### Abstract

A total forty samples of eight different medicinal plants were taken for the detection of toxigenic fungi and their mycotoxins. The fungal microflora comprises of six different fungal species belonging to three genera. *A. niger*, *Mucor species*, *A. flavus* and *Rhizopus species* dominate other fungal species isolated. Among ten samples of different medicinal plants which were contaminated with *A. flavus* was further analysed for mycotoxins potential. Four of them shows positive results for mycotoxins potential. Although the presence of toxigenic fungi in a product did not imply the presence of mycotoxins in the product, their presence represents a potential risks of contamination with mycotoxins. Therefore, these medicinal plants should be carefully stored and the growth of the naturally found toxic fungi should be inhibited. Besides that, these medicinal plants must be tested for the presence of mycotoxins present prior to their use.

**Keywords:** Medicinal plants; Microflora; Toxigenic fungi; Mycotoxins.

### Introduction

The use of Ayurveda is one of the oldest, richest, and most diverse tradition, associated with the use of medicinal plants in India (Tandon *et al.*, 2004). In this system of medicine, different parts of the medicinal plants like (bark, stem, leaves, root, fruit etc.) are used in crude as well as powered form. Over 8000 plants species have been reported to prepare some 25,000 formulation to treats the various ailments (Dubey, 2004). A large section of the Indian population unquestionably believes in the efficacy of herbal medicinal plants, and this belief has gained impetus in other countries also in the recent years. The interest in herbal medicine is due to its miraculous efficacy in curing several human ailments. Despite in the advancement in the synthetic drugs, traditional medicine system are still viable and keeping pace with the modern synthetic medicine. Popularity of this indigenous system is due to the low production cost, easy availability and fewest side effects as compare to the modern synthetic medicine.

The increasing popularity of herbal medicinal drugs made their use a Public Health problem due to lack of effective surveillance of the use, efficacy, and toxicity and quality of their natural products. The premises that traditional use of these medicinal products for generations establishes their safety does not necessarily attest to their safety and efficacy.

Indeed the adverse effects of long term herbal use, adulteration with toxic compounds and contamination by pathogenic microbial or natural toxins like mycotoxins have been reported for herbal products and medicinal plants. (Roy *et al.*, 1988; Roy and Chourasia, 1989; Efuntoye, 1999; Roy and Chourasia, 1990; Roy and Kumari, 1991; Reif, 1995; Aziz *et al.*, 1998; Halt, 1998; Abou- Arab *et al.*, 1999; Elshafie *et al.*, 1999; Freire *et al.*, 2000; Elshafie *et al.*, 2002; Rizzo *et al.*, 2004; Tassaneeyakul *et al.*, 2004 and Mandeel, 2005;).

Fungal contamination of stored herbal medicinal plants not only linked to discoloration, quality deterioration, reduction in commercial value as well in therapeutic potential but the mycotoxin produced by them in these herbal medicinal plants can also cause several ailments of liver, kidney, nervous system muscular, skin, respiratory organ, digestive tract, genital organs etc. (Purchase, 1974; Muntanola, 1987; Durakovic *et al.*, 1989; Rai and Mehrotra, 2005; Truckesses and Scott, 2008).

Quality control to prevent growth of toxigenic fungi and mycotoxins contamination in medicinal plants is essential. Keeping this in view, in the present investigation a attempt has been made to examine some stored medicinal plants samples for the association of toxigenic fungi and mycotoxins contamination under stored condition.

## Materials and Methods

### Source of samples

A total forty samples of 1. *Saraca Indica* (Ashoka) (n=5), 2. *Terminalia arjuna* (Arjuna) (n=5), 3. *Withania somnifera* (Ashwagandha) (n=5), 4. *Bacopa monnieri* (Brahmi) (n=5), 5. *Evolvulus alsinoides* (Shankhpushpi) (n=5) 6. *Zingiber officinale* (Ginger) (n=5), 7. *Tribulus terrestris* (Gokharu) (n=5) 8. *Hemidesmus indicus* (Anantmula) (n=5) were collected from the different markets of Agra and nearby regions during the year 2009 – 10. The samples were packed in air-tight polythene bags and were transported to the Department of Botany, School of Life Science, Khandari Campus, Agra, immediately from their they were stored at room temperature till further analysis.

### Measurement of moisture content

Moisture content was measured at the beginning. For moisture content, weighted amount of individual sample were dried at 100°C for 24 hours and the difference in weight before and after drying the sample was calculated according to Essono *et al.*, (2007).

$$MC = [(W_1 - W_f) / W_1] \times 100$$

where MC = moisture content,

$W_1$  = Initial weight and  $W_f$  = final weight.

### Isolation, purification and maintenance of fungal culture:

Mycoflora isolation was done by serial dilution method. The different fungi from different samples of medicinal plants was isolated and purified using standard techniques and maintenance was done on Czapek Dox Agar medium and Sabouraud's Dextrose Agar medium.

### Identification of mycoflora:

Identification of fungi was also done on the basis of morphological and cultural characteristics as described by (Barnet, 1960; Smith, 1969; Subrahmanian, 1971 and Gilman, 1975)

### Frequency of fungal species:

The Frequency of different fungal species was assessed of calculated the frequency percentage. There values were obtained according to Girridher and Ready (1997).

$$\text{Frequency percentage} = \frac{\text{No. of observation in which a species appeared}}{\text{Total no. of observations}} \times 100$$

### Preliminary screening of fungal isolate for toxigenic potential and their mycotoxins:

The different isolated fungus species obtained from the different samples of medicinal plants were tested for their Toxigenic nature by "orange, yellow" pigmentation method (Lin and Dianese, 1976).

## Results and Discussion

### Moisture content of different medicinal plant samples:

Moisture content was observed in all medicinal plants ranges form of 0.2% to 8.9%. However, it was noted that there was significant difference between the mean moisture content of different medicinal plants samples. Highest mean moisture content was observed in *E. alsinodies* (8.9%) followed by *B. monnieri* (8.7%), *Z. officinale* (7.7%), *S. indica* (3.3%), *T. arjuna* (3.2%), *W. somnifera* (2.7%), *T. terrestris* (0.3%) and the lowest mean moisture content was observed in *H. indicus* (0.2%) are shown in Table 1.

The presence of moisture content in medicinal plant samples justify favorable impact of fungal growth in stored medicinal plants (Roy, 1989; Halt, 1998). Diverse abiotic factors operating in the processing and storage conditions as well as chemicals constituents of the medicinal plants might have resulted in variation in mycopopulation in different substrates (Chourasia *et al.*, 2008). During the survey for samples collection, it was found that necessary precautions were not taken during processing and storage of these medicinal plant samples, at some places these medicinal plant samples were found under open environmental condition and some ware under dark store rooms under unhygienic conditions. All these practices may contaminate these medicinal plants by exposing them to microbial infections.

**Table 1:** Showing mean moisture content percentage in different medicinal plant samples

S.No.	Name of Plants	Mean Initial Weight $W_1$ in gm.	Mean Final Weight $W_f$ in gm.	Mean Moisture Content percentage
1.	<i>Saraca indica</i>	20	19.34	3.3
2.	<i>Terminalia arjuna</i>	20	19.36	3.2
3.	<i>Withania somnifera</i>	20	19.45	2.7
4.	<i>Bacopa monnieri</i>	20	18.25	8.7
5.	<i>Evolvulus alsinoides</i>	20	18.21	8.9
6.	<i>Zingiber officinale</i>	20	18.46	7.7
7.	<i>Tribulus terrestris</i>	20	19.94	0.3
8.	<i>Hemidesmus indicus</i>	20	19.96	0.2

**Frequency of fungal contamination:**

The mycological examination of all 40 samples of 8 different medicinal plants revealed that, 37 (92.5%) of the total analyzed samples was found to be contaminated with one or more fungal species.

All (100%) samples of *S. indica*, *T. arjuna*, *W. sominifera*, *B. monnieri*, *E. alsinoides* and *Z. officinale* were found to be contaminated with one or more fungal species.

While in *T. terrestris* 80% of the total samples analyzed was found to be contaminated with one or more fungal species.

While in *H. indicus* 60% of the total samples analyzed was found to be contaminated with one or another fungal species (Table 2)

Table : 2 Showing all the samples of different medicinal plants contaminated with one or more fungal species.

Name of Plants	Sample No.	Name of Fungus						Fungal contamination per sample
		<i>A. flavus</i>	<i>A. nidulan</i>	<i>A. niger</i>	<i>M. thermophila</i>	<i>Mucor</i>	<i>Rhizopus</i>	
<i>S. indica</i>	1	+	-	+	-	-	+	3
	2 *	+	+	-	-	-	-	2
	3	-	-	-	-	+	-	1
	4	-	-	+	-	+	-	2
	5	-	-	-	-	+	-	1
<i>T. arjuna</i>	6	+	-	+	-	-	-	2
	7	+	-	+	-	-	-	2
	8 *	+	-	+	-	-	-	2
	9	-	-	+	+	-	-	2
	10	+	-	+	-	+	-	3
<i>W. somniera</i>	11	-	-	+	-	+	-	2
	12	-	-	+	-	+	-	2
	13	-	-	+	-	+	-	2
	14	-	-	+	-	-	-	1
	15	-	-	+	-	+	-	1
<i>B.monnieri</i>	16*	+	-	+	-	-	+	3
	17	+	-	+	-	+	+	4
	18	-	-	+	-	+	+	3
	19	-	-	+	-	+	-	2
	20	-	-	+	-	-	+	2
<i>E.alsinoides</i>	21*	+	+	+	-	+	+	5
	22	-	-	+	-	-	-	1
	23	-	-	+	-	+	+	3
	24	-	-	+	-	+	+	3
	25	-	-	+	-	+	+	3
<i>Z. officinate</i>	26	-	-	+	-	+	-	2
	27	-	-	+	-	+	-	2
	28	-	-	+	-	-	+	2
	29	+	-	+	-	+	-	3
	30	-	-	+	-	+	-	2
<i>T. terrestris</i>	31	-	-	+	-	+	-	2
	32	-	-	+	-	-	-	1
	33	-	-	-	-	-	-	-
	34	-	-	-	-	+	-	1
	35	-	-	-	-	+	-	1
<i>H. indicus</i>	36	-	-	+	-	+	-	2
	37	-	-	-	-	-	-	-
	38	-	-	-	-	-	-	-
	39	-	-	-	-	+	-	1
	40	-	-	+	-	+	-	2
Total contamination in all samples		10	2	30	1	25	10	78

- Positive for mycotoxins potential.

**Distribution of different fungal species in all samples of different medicinal plants**

Analysing of different sample of all medicinal plants revealed that the frequency of contamination of *A. niger* is maximum (38.46%) followed by *Mucor* (32.05%), *Rhizopus* (12.82%), *A. flavus* (12.82%), *A. nidulans* (2.56%), and *Myceliophthora* (1.28%) as shown in Table 3.

**Table 3:** Percentage Distribution of the fungal species detected in medicinal plants.

S. No.	Fungi isolated	Number of isolates	Percentage
1.	<i>A. flavus</i>	10	12.82%
2.	<i>A. nidulans</i>	2	2.56%
3.	<i>A. niger</i>	30	38.46%
4.	<i>M. thermophila</i>	1	1.28%
5.	<i>Mucor</i> species	25	32.05%
6.	<i>Rhizopus</i> species	10	12.82%

Species of *Aspergillus* dominate the mycoflora of collected samples of medicinal plants of *S. indica*, *T. arjuna*, *W. somnifera*, *B. monnieri*, *E. alsinoides*, *G. officinale*, *T. terrestris* and *H. indicus* were it was already reported that *Aspergillus* species dominating mycoflora of stored medicinal plants (Hitokoto *et al.*, 1978; Ayres *et al.*, 1980; Aziz *et al.*, 1998; Arab *et al.*, 1999; Elshafie *et al.*, 1999; Mandeel, 2005). Hence, the presence of wide range of fungi in these medicinally important medicinal plants showed that there was is a potential risk for mycotoxins contamination, especially during prolonged storage in poor storing conditions without temperature and moisture control (Efuntoyc 2004; Bugno *et al.*, 2006; Singh *et al.*, 2008).

**Distribution of different fungal species with in samples of different medicinal plants:**

By Analyzing the samples of different medicinal plants it was observed that the samples of *S. indica* medicinal plant samples were frequently contaminated with *Aspergillus flavus* (40%) *A. nidulans* (20%), *A. niger* (40%), *Mucor* (60%) and *Rhizopus* (20%), in *T. arjuna* medicinal plant samples *A. flavus* (80%), *A. niger* (100%), *M. thermophila* (20%), *Mucor* (20%), in *W. sominifera* medicinal plant samples *A. niger* (80%), *Mucor* (80%), in *B. monnieri* samples *A. flavous* (40%), *A. niger* (100%), *Mucor* (60%), *Rhizopus* (80%), in *E. alsinoides* samples *A. flavus* (20%) *A. nidulans* (20%), *A. niger* (100%), *Mucor* (80%), *Rhizopus* (80%), in *Z. officinate* samples *A. flavus* (20%), *A. niger* (100%), *Mucor* (80%), *Rhizopus* (20%), in *T. terrestris* samples *A. niger* (40%), *Mucor* (60%) and in *H. indicus* samples *A. niger* (40%), *Mucor* (60%) were found (Table 4.)

**Determination of toxigenic contamination of samples**

It is known that, some fungi can synthesize orally toxic metabolites – mycotoxins. So, in our present study we paid our attention to study the potential of *A. flavus* fungal species to produced mycotoxin. The isolate of *A. flavus* was subcultured on mycotoxin activator media (Coconut Agar Media). According to the results out of total ten samples of different medicinal plants which was contaminated with *A. flavus* were subcultured on mycotoxin activating media, four samples are found as mycotoxins producing fungi (Table 5).

Fungal contamination of raw materials of medicinal plants is a major impediment preventing India from becoming a herbal giant. Therefore, fungal contamination in medicinal plants, especially raw materials, should be prevented during storage. Plant materials used for medical purposes should be carefully stored and the growth of toxigenic fungi should be inhibited. In the present study 40 samples of eight different medicinal plants were collected from different markets places of Agra and nearby regions were taken to evaluate the presence of toxigenic fungi. Out of 40 samples of different medicinal plants 37 (92.57%) samples are found to be contaminated with one or more fungal species. All (100%) the samples of *S. indica*, *T. arjuna*, *W. sominifera*, *B. monnieri*, *E. alsinoides* and *Z. officinale* were found to be contaminated with one or more fungal species, while 80% in the case of *T. terrestris* and 60% in case of *H. indicus* was found to be contaminated with one or more fungal species

**Table 4:** Percentage of samples contamination by different fungal species isolated from different medicinal plants.

Fungal Species	% of samples yielding different species of fungi							
	<i>S. indica</i>	<i>T. arjuna</i>	<i>W. somnifera</i>	<i>B. monnieri</i>	<i>E.alsinoides</i>	<i>Z.officinale</i>	<i>T. terrestris</i>	<i>H. indicus</i>
<i>A. flavus</i>	40%	80%	-	40%	20%	20%	-	-
<i>A. nidulans</i>	20%	-	-	-	20%	-	-	-
<i>A. niger</i>	40%	100%	80%	100%	100%	100%	40%	40%
<i>M. thermophila</i>	-	20%	-	-	-	-	-	-
<i>Mucor</i> species	60%	20%	80%	60%	80%	80%	60%	60%
<i>Rhizopus</i> species	20%	-	-	80%	80%	20%	-	-

**Table 5:** Distribution of mycotoxin producing *A. flavus* among different medicinal plant samples

S. No.	Medicinal plant sample	Total number of fungi examined	Positive strains of mycotoxin production
1.	<i>S. indica</i>	2	1
2.	<i>T. arjuna</i>	4	1
3.	<i>W. somnifera</i>	-	-
4.	<i>B. monnieri</i>	2	1
5.	<i>E. alsinoides</i>	1	1
6.	<i>Z. officinale</i>	1	-
7.	<i>T. terrestris</i>	-	-
8.	<i>H. indicus</i>	-	-
	Total	10	4

(-) not detected

The mycoflora comprise of six different fungal species belonging to three genera viz. The maximum fungal species are of *A. niger* (38.46%), followed by *Mucor* species (32.05%), *Rhizopus* species (12.82%), *A. flavus* (12.82%), *A. nidulans* (2.56%) and *M. thermophila* (1.82%). Among fungi isolated the presence of *A. niger* 38.46%, *Mucor* 32.05%, *A. flavus* 12.82% and *Rhizopus* species 12.82% dominate other fungal species isolated. Among total 10 samples of different medicinal plants which were found to be contaminated with *A. flavus* were further analyzed for mycotoxin potential by subcultured on mycotoxin activating media (coconut agar media) 4 out of 10 shows positive toxigenic potential for mycotoxins.

## References

Abou-Arab AAK, Kawther MS, El Tantawy ME, Badaea RI and Khayria N (1999) Quantity estimation of some contaminants in commonly used medicinal plants in the

Egyptian market. *Food Chem.* **67**: 357-363. DOI: 10.1016/S0308-8146(99)00082-5

Aryes GI, Mund TI and Sondin EW (1980) Microbiology of food spices and condiments. A series of books in food and nutrition Edn. Schmeigert, pp 249.

Aziz NH, Youssef YA, Moheye ZEL-F and Lofty AM (1998) Contamination of some common medicinal plant samples and spices by fungi and their mycotoxins. *Botanical Bulletin of Academic Sciences* **39**: 279-285.

Barnet UL (1960) Illustrate genera of imperfect fungi. *1<sup>st</sup> ed. Minneapolis Burgess Publ. Co.*

Bugno A, Adriana ABA, Tatiana CP, Terezinha, AP and Myrna S (2006) Occurrence of toxigenic fungi in herbal drugs. *Brazilian Journal of Microbiology* **37**: 47-51. DOI: 10.1590/S1517-83822006000100009

Chourasia HK, Suman SK and Prasad P (2008) Mycotoxic fungi and mycotoxins in fast food of Bihar. *Journal of Mycology and Plant Pathology* **38**: 492-499.

Dubey NK, Kumar R and Tripathi P (2004) Global promotion of herbal medicine: India's opportunity. *Current Science* **86**: 37- 41. DOI: 10.1016/j.lfs.2005.09.012.

Durakovic S, Galic J and Pajnovic P (1989) Toxic and cancer metabolites of moulds in food and fodder. *Hrana Iishrana* **30**: 71-100.

Efuntoye MO (1999) Mycotoxins of fungal strains from stored herbal plants and mycotoxin contents of Nigerian crude herbal drugs. *Mycopathologia.* **147**: 43-48. DOI: 10.1023/A:1007007418581

Efuntoye MO (2004) Fungi associated with herbal drug plants during storage. *Mycopathologia.* **136**: 115-118. DOI: 10.1007/BF00437505

Elshafie AE, Al-Lawatia T and Al- Bahry S (1999) Fungi associated with black tea and tea quality in the Sultanate of Oman. *Mycopathologia* **145**: 89-93. DOI: 10.1023/A:1007034729467

Elshafie AE, Al-Rashdi, TA, Al-Bahry SN and Bakheit CS (2002) Fungi and aflatoxins associated with spices in the Sultanate of Oman. *Mycopathologia.* **155**: 155-160. DOI: 10.1023/A:1020427527963

Essono G, Ayodele M, Akoa A, Foko J, Olembo S and Gock wski J (2007) *Aspergillus* species on cassava chips in storage in

- rural areas of Southern Cameroon: their relationship with storage duration, moisture content and processing methods. *Afric. J. Microbiol.* **1**: 1-8.
- Gilman JC (1975) A manual of soil fungi. *Oxford and IBH Publishing Corporation, New Delhi, Bombay, Calcutta.*
- Giridher P and Ready SM (1997) Incidences of mycotoxin producers on spices from Andhra Pradesh, *J. Indian Bot. Soc.* **76**: 161-164.
- Halt M (1998) Moulds and mycotoxins in herb tea and medicinal plants, *European Journal of Epidemiology* **14**: 269–274. DOI: 10.1023/A:1007498613538
- Hitokoto H, Morozumi S, Wauke T, Saka S and Kurata H (1978) Fungal contamination and mycotoxin detection of powdered herbal drugs. *Applied and Environmental Microbiology* **36**: 252-256.
- Lin MT and Dlanese JC (1976) A coconut agar medium for rapid detection of mycotoxins produced by *Aspergillus* Spp. *Phytopathol.* **66**: 1466-1468.
- Mandee QA (2005) Fungal contamination of some imported spices. *Mycopathologia.* **159**: 291-298. DOI: 10.1007/s11046-004-5496-z
- Muntanola M (1987) General mycology. Beograd: NIRO. *Knjez evne novine* 257-269.
- Purchase IFH (1974) Mycotoxins. Amsterdam. *Elsevier*, 1–28.
- Rai V and Mehrotra S (2005) Toxic Contaminants in Herbal Drugs. *Environ. News* **11**: 1-3.
- Reif K and Metzger W (1995) Determination of aflatoxins in medicinal herbs and plant extracts. *J. Chromatography A.* **692**: 131-136. DOI: 10.1016/0021-9673(94)00841-V
- Rizzo I, Vedoya G, Maurutto S, Haidukowski M and Varsavsky E (2004) Assessment of toxigenic fungi on Argentinean medicinal herbs. *Microbiol. Res.* **159**(2): 113-120. DOI: 10.1016/j.micres.2004.01.013
- Roy AK, Sinha KK and Chourasia HK (1988) Aflatoxin contamination of some common drug plants. *Appl. Environ. Microbiol.* **54**: 842-843.
- Roy AK and Chourasia HK (1989) Aflatoxin problem in some medicinal plants under storage. *Int. Jour. Crude Drug Res.* **27**: 156-160.
- Roy AK and Chourasia HK (1990) Effect of relative humidity on fungal association and aflatoxin production on Piper longum fruits. *Ind. Bot. Repr.* **8**: 138-140.
- Roy AK and Kumari V (1991) Aflatoxin and citrinin in seeds of some medicinal plants under storage. *Int. J. Pharmacognosy.* **29**: 62-65. DOI: 10.3109/13880209109082851
- Singh P, Srivastava B, Kumar A and Dubey NK (2008) Fungal Contamination of Raw Materials of Some Herbal Drugs and Recommendation of Cinnamomum camphora Oil as Herbal Fungitoxicant. *Microbial Ecology.* **56**(3): 555-560 DOI: 10.1007/s00248-008-9375-x
- Smith G (1969) An introduction to industrial mycology. *Eward Arnold Ltd. London.* 390.
- Soares LMV and Aodriguez-Amaya DB (1989) Survey of aflatoxins, ochratoxins A, zearalenone and sterigmatocystin in some Brazilian foods by using multi-toxin thin layer chromatographic method. *J. Assoc. Off. Anal. Chem.* **72**: 22 – 26.
- Subrahmanian CV (1971) 'Hyphomycetes' ICAR, New Delhi. 1930.
- Tassaneeyakul W, Razzazi-Fazeli E, Porasuphatana S and Bohm J (2004) Contamination of aflatoxins in herbal medicinal products in Thailand. *Mycopathologia* **158**: 239-244. DOI: 10.1023/B:MYCO.0000041892.26907.b4
- Truckesses MW and Scott PM (2008) Mycotoxins in botanicals and dried fruits: A review. *Food Additives and Contaminants* **25**: 181–192. DOI: 10.1080/02652030701567459