



# International Journal of Applied Sciences and Biotechnology

A Rapid Publishing Journal

ISSN 2091-2609

## Indexing and Abstracting

CrossRef, Google Scholar, Global Impact Factor, Genamics, Index Copernicus, Directory of Open Access Journals, WorldCat, Electronic Journals Library (EZB), Universitätsbibliothek Leipzig, Hamburg University, UTS (University of Technology, Sydney): Library, International Society of Universal Research in Sciences (EyeSource), Journal Seeker, WZB, Socolar, BioRes, Indian Science, Jadoun Science, Jour-Informatics, Journal Directory, JournalTOCs, Academic Journals Database, Journal Quality Evaluation Report, PDOAJ, Science Central, Journal Impact Factor, NewJour, Open Science Directory, Directory of Research Journals Indexing, Open Access Library, International Impact Factor Services, SciSeek, Cabell's Directories, Scientific Indexing Services, CiteFactor, UniSA Library, InfoBase Index, Infomine, Getinfo, Open Academic Journals Index, HINARI, etc.

**CODEN (Chemical Abstract Services, USA): IJASKD**

Vol-3(4) December, 2015

Available online at:

<http://www.ijasbt.org>

&

<http://www.nepjol.info/index.php/IJASBT/index>



Impact factor\*: **1.422**

Scientific Journal Impact factor#: **3.419**

Index Copernicus Value: **6.02**

IBI Factor 2015\*\*: **4.19**

\*Impact factor is issued by Universal Impact Factor. Kindly note that this is not the IF of Journal Citation Report (JCR).

For any type of query and/or feedback don't hesitate to email us at: [editor.ijasbt@gmail.com](mailto:editor.ijasbt@gmail.com)



Research Article

## AZOLLA AS AN ECONOMIC SUBSTITUTE TO SOYBEAN BASED FEED FOR POULTRY

Dev R. Paudel<sup>1</sup>, Pramila Dhakal<sup>2</sup>, Kailash P. Timsina<sup>1</sup> and Ananta Dahal<sup>1</sup>

<sup>1</sup>Institute of Agriculture and Animal Science, Rampur, Chitwan, Nepal

<sup>2</sup>Food and Agriculture Organization of the United Nations, Surkhet, Nepal

Corresponding author email: dpaudel@outlook.com

### Abstract

A study on azolla as an economical substitute to soybean based poultry feed was conducted. The experiment consisted of four treatments laid out in Completely Randomized Design (CRD) with five replications. Each experimental unit consisted of three birds. For this purpose, eight-day old, unsexed Hubbard broiler chicks were weighed and randomly assigned to each treatment. Four different levels of azolla meal were incorporated in the starter and finisher ration at a level of 0%, 10%, 15%, and 20% respectively. Body weight, weight gain, feed consumption, and feed efficiency were measured. The average cumulative weekly body weight for all weeks were found to be significantly different ( $P < 0.05$ ) for the various treatments. Maximum cumulative body weight (1360 g) was found in broiler fed T2 ration in 5th week. Similar trend was also observed in 6th week showing highest cumulative weight (1882 g) in control diet (T1) which was at par with T2 (1841 g). Significantly highest (420.3 g) live weight gain was observed in T2 diet during 5th week. Overall mean of weekly feed consumption was recorded highest (784.33 g) in T2 diet. The economic analysis of net income/kg live weight was the highest (Rs 18.86) in T1 which was at par with T2 (Rs 18.35). Likewise, cost of production/bird was highest (Rs 124.5) in T1 and lowest (Rs 112.2) in T4. The results show that azolla meal could be utilized to substitute up to 10% soybean meal from overall production and economic point of view.

**Keywords:** Azolla; Poultry; Feed; Sustainable; Protein; Substitute; Economic

### Introduction

Poultry farming has emerged as a major income generating enterprise in agriculture sector within the reach of the poor, women, marginal farmers, and entrepreneurs in Nepal. It has been playing a significant role in the rural economy and its contribution to national GDP and AGDP is estimated to be around 3% and 10% respectively involving more than 60,000 families (USAID, 2008).

Feed alone costs 65 to 75% of the total broiler production and ultimately results in high price of poultry meat (Johari and Hussain, 1996). Chicken has very limited ability to meet their protein needs from non-protein sources. Typical broiler ration will contain 22 to 24% protein. Protein synthesis in chickens is required at a very rapid rate for compensating the broken tissues of the adult body (Banerjee, 2000). The feeding of poultry is done through concentrates. Commercial feeds are expensive for average Nepalese farmers as more than 55 percent of Nepal's population lives below the international poverty line of \$1.25 per day (FTF, 2010). Therefore, the poultry rearing farmers are in search of economical and sustainable feed ingredients for poultry feeding. Many researchers are working on different materials to be used for economic broiler feed ration formulation (Olorunfemi et al., 2006;

Fasuyi et al., 2007; Agbede, 2003; Abeke et al., 2008). Azolla has shown a great potential to be included as a source of protein for duck ration (Bacerra et al., 1995) and egg type chicks (Alalade and Iyayi, 2006).

Azolla is a floating fern very rich in protein, essential amino acids, vitamins (Vit. A, Vit. B12, Beta carotene), growth promoter intermediaries and minerals like calcium, phosphorus, potassium, ferrous, copper, magnesium, etc. (Kamalasanana et al., 2002). A blue-green algae (*Anabaena azollae*) lives in the cavities of azolla leaves and fixes nitrogen from the atmosphere. Azolla has been used for different agricultural purposes (Akhter et al., 2002; Krishnakumar et al., 2005; Hossain et al., 2001; Tahmid et al. 2002). Azolla has been used in animal feed ration as well (Sheeno and Sahu, 2006). The bio-composition of azolla makes it one of the most economic, efficient and sustainable feed substitute for poultry (Kamalasanana et al., 2002). Incorporation of azolla as an alternative protein ingredient in poultry ration could make poultry production economical.

## Materials and Method

### Experimental Site

The experiment was conducted at livestock farm of Institute of Agriculture and Animal Science (IAAS), Rampur, Chitwan.

### Husbandry/Experimental Design

Sixty, day old Hubbard chicks were obtained from commercial hatchery and reared in battery brooder for one week. After brooding, chicks were transferred to experimental pen. The experiment was conducted in Completely Randomized Design (CRD) consisting of four treatments along with five replications. Each experimental unit consisted of 3 birds.

### Azolla Meal Preparation

Whole plants of *Azolla pinnata* were harvested in sufficient quantities from Rupa Lake, Pokhara; Sangle river, Kathmandu and IAAS fish pond. The collected azolla was sun dried for three days until they became crispy while retaining greenish coloration. The dried leaves were then milled and used as feed ingredient. Chemical composition of the azolla was analyzed and is presented in Table 1. The analysis was carried out according to the standard methods of A.O.A.C. (1990).

**Table 1:** Chemical composition of Azolla

Constituents	Percentage
Dry matter	89.1
Crude protein	24.18
Crude fiber	15.41
Ether extract	3.35
Nitrogen free extract	30.04
Total ash	16.12

### Ration Formulation

Two types of rations, one ration for broiler starter (0-4 weeks) and another ration for broiler finisher (4-6 weeks) were formulated (Table 2 and Table 3).

### Feeding Trial

4 treatments with varying levels of *Azolla* meal were prepared as follows:

Symbol	Treatment details
T1	Azolla meal AZM 0%
T2	Azolla meal AZM 10%
T3	Azolla meal AZM 15%
T4	Azolla meal AZM 20%

**Table 2:** Details of treatment composition (ingredients) and nutrient composition used in broiler starter rations (0-4 weeks)

Ingredients	Treatments			
	T <sub>1</sub>	T <sub>2</sub>	T <sub>3</sub>	T <sub>4</sub>
Maize	582	544	526	529
Rice polish	50	60	53	22
Azolla meal	0	100	150	200
Soybean meal	310	256	232	216
Bone meal	20	20	20	20
Marble	30	15	14	10
Common salt	2	2	2	2
Vit.-Min. Premix	+	+	+	+
Calculated nutrient composition				
ME Kcal/kg	2886.8	2840.3	2802.75	2775.8
CP%	20.788	20.276	20.020	20.025
CF%	3.748	4.658	5.039	5.327
Ca%	1.721	1.284	1.298	1.205
Av. P%	0.671	0.766	0.800	0.815

**Table 3:** Details of treatment composition (ingredients) and nutrient composition used in broiler finisher rations (5-6 weeks)

Ingredients	Treatments			
	T <sub>1</sub>	T <sub>2</sub>	T <sub>3</sub>	T <sub>4</sub>
Maize	625	646	620	598
Rice polish	100	22	20	10
Azolla meal	0	100	150	200
Soybean meal	250	220	199	181
Bone meal	10	5	5	5
Marble	10	5	4	4
Common salt	2	2	2	2
Vit.-Min. Premix	+	+	+	+
Calculated nutrient composition				
ME Kcal/kg	3015.50	2990.33	2949.85	2905.73
CP%	18.149	18.036	18.074	18.019
CF%	2.897	3.228	3.753	4.196
Ca%	0.725	0.517	0.531	0.581
Av. P%	0.604	0.548	0.589	0.621

**General Management**

Available brooder was washed, cleaned, and disinfected using Kohrsolin @ 0.03% solution. Floors, interior of the walls and roof were scrubbed and left over litters were removed from the brooder house. The poultry house was thoroughly cleaned. Before placing the litter in poultry house, dust lime was applied as disinfectant. In all experimental pens, rice husk was used as bedding material. The thickness of litter on floor was made to 10 cm. The birds were fed *ad libitum* in all treatments up to the 6th week of age.

**Data Collection****Weekly body weight and body weight gain**

The initial and weekly body weights of the birds were taken. Body weight gain was obtained by subtracting previous weight of the birds from their corresponding body weight for each week.

**Feed and Water Consumption**

The chicks of different treatments were reared under similar management and environmental conditions. Feed was offered *ad libitum* every day during the entire experimental period. Weekly ration consumption of each treatment was recorded by subtracting the weight of the left over ration from the weight of ration offered in each week. Water pots were thoroughly cleaned before watering each time. Fresh water was made available *ad libitum* during the experimental period.

**Feed Conversion Ratio**

The weekly feed conversion ratio of birds in each treatment was determined by dividing the weekly feed intake by their respective weight gain.

Feed Conversion Ratio (FCR) = Weekly feed intake (g) / Weekly weight gain (g)

**Economics of Production**

The economics of broiler production was calculated at the end of the experiment. Gross expenditure was calculated by adding the cost of chicks, feeds, medicines, management, and labor cost. The gross income was obtained by adding selling price of the birds (NRs 75/bird) [USD 1 = Rs. 75] and litter (Rs 30/bag) at local market. The different economic parameters were calculated as follows.

Net income per bird (Rs.) = [Gross income (Rs.) – Gross expenditure (Rs.)] / No. of live birds

Net income per kg live weight (Rs.) = Net income of birds (Rs.) / Total live weight of birds (kg)

Cost of production per bird (Rs.) = Gross expenditure (Rs.) / Total number of birds in treatment

**Statistical Analysis**

The data recorded on different parameters during experiment were first tabulated in Microsoft Excel and then Analysis of Variance (ANOVA) for all data was statistically analyzed using MSTAT-C computer software package.

**Results and Discussion****Feed Consumption**

The analysis of variance showed a non-significant difference ( $P>0.05$ ) among the treatments during the second and third week of the experimental period (Table 4). A significant difference in the average weekly feed consumption was observed in the 4th week. The highest feed consumption (850.0 g) was found in T4 which was at par with T2 (836.0 g) and T1 (799.3 g). The lowest feed consumption was found in T3 (701.7 g). Likewise in the fifth week, significantly highest feed consumption (1031.0 g) was found in T2 which was at par with T3 (971.3 g) and the lowest feed consumption was found in T4 (834.3 g) which was at par with T1 (856.0 g). In the 6th week, significantly highest feed consumption was found in T2 (1202.0 g) which was at par with T1 (1182.0 g) and T3 (1111.0 g). The lowest feed consumption was found in T4 (974.0 g).

**Body Weight**

A significant difference ( $P<0.05$ ) between the treatments was observed in all weeks for body weight (Table 5). During the 6th week, significantly highest body weight was found in T1 (1882.0 g) which was at par with T2 (1841.0) and the lowest body weight was found in T4 (1517 g).

The results are similar to Cambel (1984) who found better result using 10% and 15% AZM. The higher level of azolla (T3 and T4) meal resulted in poor growth of the broilers as compared to the growth of the broilers in T1 and T2 treatments.

**Weekly Weight Gain**

Significantly highest weekly weight gain (137.7 g) in 2nd week was recorded in T1 and the lowest weight gain (117.7 g) was recorded in T4 which was at par with T2 (122.3 g) and T3 (119.0 g) (Table 6). In the 6th week, significantly highest weight gain was observed in T1 (525.3 g) and the lowest weight gain was observed in T4 (362.0 g).

The above results are similar to Cambel (1984) who found highest weekly weight gain in 10% AZM dietary treatment. Alalade and Iyayi, 2006 also reported that diet containing 10% AZM did non-significantly better than the control.

**Feed Conversion Ratio (FCR)**

A significant difference ( $P<0.05$ ) in the FCR was observed in all weeks (Table 7). In the sixth week poorer FCR (2.690) was observed in T4 which was at par with T3 (2.620) and superior FCR (2.252) was observed in T1.

**Table 4:** Average weekly feed consumption (g) of Hubbard broilers fed with diets using azolla meal as a feed ingredient in broiler ration

Treatment	Period (week) and feed consumption (g)					Overall mean
	2	3	4	5	6	
T <sub>1</sub> = AZM 0%	290.7	577.9	799.3 <sup>a</sup>	856.0 <sup>b</sup>	1182.0 <sup>a</sup>	741.18
T <sub>2</sub> = AZM 10%	285.3	569.3	836.0 <sup>a</sup>	1031.0 <sup>a</sup>	1202.0 <sup>a</sup>	784.72
T <sub>3</sub> = AZM 15%	291.0	567.7	701.7 <sup>b</sup>	971.3 <sup>a</sup>	1111.0 <sup>a</sup>	728.54
T <sub>4</sub> = AZM 20%	295.3	629.0	850.0 <sup>a</sup>	834.3 <sup>b</sup>	974.0 <sup>b</sup>	716.52
Mean	290.6	586.0	779.2	921.3	1139.7	
Probability	0.942	0.1898	0.0104	0.0000	0.0027	
LSD	34.38	65.07	87.64	65.01	114.5	
SEm <sub>±</sub>	11.47	21.70	29.23	21.68	38.21	
CV%	8.83	8.28	8.20	5.25	7.65	

Means within the column followed by different superscripts are significantly different (P<0.05) by LSD

**Table 5:** Average cumulative weight (g) of Hubbard broilers fed with diets using azolla meal as a feed ingredient in broiler ration

Treatment	Period (week) and feed consumption (g)					
	Initial wt.	2	3	4	5	6
T <sub>1</sub> = AZM 0%	159.0 <sup>a</sup>	296.7 <sup>a</sup>	587.7 <sup>a</sup>	973.3 <sup>a</sup>	1356.7 <sup>a</sup>	1882.0 <sup>a</sup>
T <sub>2</sub> = AZM 10%	150.3 <sup>b</sup>	272.7 <sup>b</sup>	545.3 <sup>b</sup>	940.0 <sup>a</sup>	1360.3 <sup>a</sup>	1840.7 <sup>a</sup>
T <sub>3</sub> = AZM 15%	148.7 <sup>b</sup>	267.7 <sup>b</sup>	520.0 <sup>b</sup>	837.7 <sup>b</sup>	1224.7 <sup>b</sup>	1648.7 <sup>b</sup>
T <sub>4</sub> = AZM 20%	146.0 <sup>b</sup>	263.7 <sup>b</sup>	516.0 <sup>b</sup>	840.3 <sup>b</sup>	1155.3 <sup>c</sup>	1517.3 <sup>c</sup>
Mean	151.0	275.2	542.25	897.8	1274.25	1722.17
Probability	0.0011	0.0029	0.0003	0.0000	0.0000	0.0000
LSD	5.699	16.60	29.42	43.51	55.91	75.37
SEm <sub>±</sub>	1.901	5.538	9.813	14.51	18.65	25.14
CV%	2.82	4.50	4.05	3.61	3.27	3.26

Means within the column followed by different superscripts are significantly different (P<0.05) by LSD

**Table 6:** Average weekly body weight gain (g) of Hubbard broilers fed with diets using azolla meal as a feed ingredient in broiler ration

Treatment	Period (week) and feed consumption (g)					Overall mean
	2	3	4	5	6	
T <sub>1</sub> = AZM 0%	137.7 <sup>a</sup>	291.0 <sup>a</sup>	385.7 <sup>a</sup>	383.3 <sup>b</sup>	525.3 <sup>a</sup>	344.6
T <sub>2</sub> = AZM 10%	122.3 <sup>b</sup>	272.7 <sup>a</sup>	394.7 <sup>a</sup>	420.3 <sup>a</sup>	480.3 <sup>b</sup>	338.1
T <sub>3</sub> = AZM 15%	119.0 <sup>b</sup>	252.3 <sup>b</sup>	317.7 <sup>b</sup>	387.0 <sup>b</sup>	424.0 <sup>c</sup>	300.0
T <sub>4</sub> = AZM 20%	117.7 <sup>b</sup>	252.3 <sup>b</sup>	324.3 <sup>b</sup>	315.0 <sup>c</sup>	362.0 <sup>d</sup>	274.3
Mean	124.17	267.07	355.6	376.4	447.9	
Probability	0.0306	0.0012	0.0000	0.0000	0.0000	
LSD	14.12	19.00	27.99	21.34	41.69	
SEm <sub>±</sub>	4.710	6.337	9.336	7.119	13.91	
CV%	8.48	5.31	5.87	4.23	6.94	

Means within the column followed by different superscripts are significantly different (P<0.05) by LSD



**Table 7:** Average feed conversion ratio of Hubbard broilers fed with diets using azolla meal as a feed ingredient in broiler ration

Treatment	Period (week) and feed consumption (g)					Overall mean
	2	3	4	5	6	
T <sub>1</sub> = AZM 0%	2.110 <sup>c</sup>	1.984 <sup>c</sup>	2.072 <sup>c</sup>	2.236 <sup>c</sup>	2.252 <sup>c</sup>	2.13
T <sub>2</sub> = AZM 10%	2.336 <sup>b</sup>	2.090 <sup>c</sup>	2.114 <sup>c</sup>	2.456 <sup>b</sup>	2.5 <sup>b</sup>	2.30
T <sub>3</sub> = AZM 15%	2.446 <sup>ab</sup>	2.250 <sup>b</sup>	2.210 <sup>b</sup>	2.510 <sup>b</sup>	2.620 <sup>a</sup>	2.41
T <sub>4</sub> = AZM 20%	2.510 <sup>a</sup>	2.490 <sup>a</sup>	2.620 <sup>a</sup>	2.650 <sup>a</sup>	6.690 <sup>a</sup>	2.59
Mean	2.35	2.20	2.26	2.46	2.52	
Probability	0.0001	0.0000	0.0000	0.0001	0.0000	
LSD	0.1469	0.1406	0.1122	0.1406	0.09481	
SEm <sub>±</sub>	0.04899	0.04690	0.03742	0.04690	0.03162	
CV%	4.62	4.84	3.72	4.29	2.77	

Means within the column followed by different superscripts are significantly different ( $P < 0.05$ ) by LSD

The above results are similar to Basak et al. (2002) who reported poorest feed conversion ratio in 15% AZM included diets. This might be due to higher fiber content of azolla. Higher level of fiber and tannin in aquatic plant may be responsible for decreasing the nutrient utilization and ultimately decrease FCR (Muzlar et al., 1978).

#### Survivability

None of the birds died in any treatments during the experimental period. This indicates that azolla meal had no deleterious effects on broilers. The results are similar with Basak et al. (2002) and Castilo et al. (1981) who also found no toxic effect of dietary azolla.

#### Economics of Broiler Production

The analysis of variance was statistically significant ( $P < 0.05$ ) for all the economic parameters (Table 8). The cost of production/bird was highest (Rs124.5) in T<sub>1</sub> which was at par with T<sub>2</sub> (Rs 122.6) and the lowest cost of production/bird (Rs 112.2) was recorded in T<sub>4</sub> which was at par with T<sub>3</sub> (Rs 115.3). Similarly, the cost of production/kg live weight (Rs 66.14) was recorded in T<sub>1</sub> which was at par with T<sub>2</sub> (Rs 66.65). Likewise, highest gross income/bird was found in T<sub>1</sub> (Rs 160.0) which was at par with T<sub>2</sub> (Rs 156.5) and the lowest gross income/bird was found in T<sub>4</sub> (Rs 129.0).

Net income/bird was observed highest in T<sub>1</sub> (Rs 35.50) which was at par with T<sub>2</sub> (Rs 33.88) and the lowest net income/bird was observed in T<sub>4</sub> (Rs 16.80). Likewise, the net income/kg live weight was found highest (Rs 18.86) in T<sub>1</sub> which was at par with T<sub>2</sub> (Rs 18.35) and the lowest net

income/kg live weight was in T<sub>4</sub> (Rs 11.02). The percentage net income was recorded maximum (28.54%) in T<sub>1</sub> which was at par with T<sub>2</sub> (27.59%) and minimum on T<sub>4</sub> (14.93%).

#### Conclusion

The use of azolla meal as a feed ingredient in broiler ration was investigated. The experiment consisted of four treatments of azolla meal at 0%, 10%, 15% and 20%, laid out in the completely randomized design (CRD) with five replications. Each experimental unit consisted of three birds. Two types of rations, broiler starter (0 to 4th week) and finisher (5th to 6th week) were fed. The highest average total feed consumption from 2nd to 6th weeks of age was obtained on T<sub>2</sub> (3923.6 g) and the lowest on T<sub>4</sub> (3582.6 g). The highest final average body weight was obtained on T<sub>1</sub> (1882 g) which was at par with T<sub>2</sub> (1841 g) and the lowest body weight was found on T<sub>4</sub> (1517.7 g). The average weekly body weight gain of broilers observed was statistically significant ( $P < 0.05$ ) for all weeks except 2nd week. The average weekly feed conversion ratio was the best (2.13) in control (T<sub>1</sub>) diet and was at par with T<sub>2</sub> (2.30) and poorest in T<sub>4</sub> (2.59). Overall economic analysis of broiler production was observed to be significantly different ( $P < 0.05$ ) among the treatments.

The inclusion of azolla meal at 10% (T<sub>2</sub>) was either superior or at par with control diet (T<sub>1</sub>) with respect to overall performance and net economic returns. From the results it can be concluded that broilers could be raised successfully without compromising performance while substituting soybean meal up to 10% with azolla meal.

**Table 8:** Cost of production and net returns at different dietary treatments

Treatment	Gross expenditure (Rs)	Cost of production/bird (Rs)	Cost of production/kg live weight (Rs)	Gross income (Rs)	Gross income/bird (Rs)	Net income (Rs)	Net income/bird (Rs)	Net income/kg live weight (Rs)	Net income (%)
T <sub>1</sub> = AZM 0%	373.4 <sup>a</sup>	124.5 <sup>a</sup>	66.14 <sup>c</sup>	479.9 <sup>a</sup>	160.0 <sup>a</sup>	106.5 <sup>a</sup>	35.50 <sup>a</sup>	18.86 <sup>a</sup>	28.54 <sup>a</sup>
T <sub>2</sub> = AZM 10%	367.7 <sup>a</sup>	122.6 <sup>a</sup>	66.65 <sup>c</sup>	469.4 <sup>a</sup>	156.5 <sup>a</sup>	101.6 <sup>a</sup>	33.88 <sup>a</sup>	18.35 <sup>a</sup>	27.59 <sup>a</sup>
T <sub>3</sub> = AZM 15%	345.9 <sup>b</sup>	115.3 <sup>b</sup>	69.93 <sup>b</sup>	420.4 <sup>b</sup>	140.1 <sup>b</sup>	74.55 <sup>b</sup>	24.85 <sup>b</sup>	15.07 <sup>b</sup>	21.56 <sup>b</sup>
T <sub>4</sub> = AZM 20%	336.5 <sup>b</sup>	112.2 <sup>b</sup>	73.98 <sup>a</sup>	386.9 <sup>c</sup>	129.0 <sup>c</sup>	50.39 <sup>c</sup>	16.80 <sup>c</sup>	11.02 <sup>c</sup>	14.93 <sup>c</sup>
Mean	355.9	118.6	69.2	439.2	146.4	83.3	27.8	15.8	23.1
Probability	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000
LSD	10.35	3.451	1.609	19.22	6.406	11.22	3.740	1.609	2.855
SEm±	3.453	1.151	0.5367	6.410	2.137	3.742	1.247	0.5367	0.9524
CV%	2.17	2.17	1.73	3.26	3.26	10.05	10.05	7.58	9.20

Means within the column followed by different superscripts are significantly different (P<0.05) by LSD

## Acknowledgements

The authors would like to thank Global Future Institute (GFI) Small Research Grant Program, USA for the financial support provided for this research, Dr. Ashesh Bhattarai for feed formulation, and Dr. Ishwor Chandra Prakash Tiwari for facility and laboratory use.

## References

- Abeke FO, Ogundipe SO, Sekoni AA, Adeyinka IA, Oni OO, Abeke A, and Dafwang II (2008) Effect of Dietary Levels of Cooked Lablab purpureus Beans on the Performance of Broiler Chickens. *American Journal of Food Technology*. **3**: 42-49. DOI: 10.3923/ajft.2008.42.49
- Agbede JO (2003) Equi-protein Replacement of Fishmeal with Leucaena Leaf Protein Concentrate: An Assessment of Performance Characteristics and Muscle Development in the Chicken. *International Journal of Poultry Science*. **2**: 421-429. DOI: 10.3923/ijps.2003.421.429
- Akhter S, Mian MH, Kader MA, and Begum SA (2002) Combination of Azolla and Urea Nitrogen for Satisfactory Production of Irrigated Boro Rice (BRRI Dhan 29). *Journal of Agronomy*. **1**: 127-130. DOI: 10.3923/ja.2002.127.130
- Alalade OA and Iyayi EA (2006) Chemical Composition and the Feeding Value of Azolla (*Azolla pinnata*) Meal for Egg-Type Chicks. *International Journal of Poultry Science* **5** (2): 137-141. DOI: 10.3923/ijps.2006.137.141
- AOAC (1990) Official method of analysis, Association of official Analytical Chemists (15th ed.), Arlington, VA.
- Bacerra M, Preston TR and Ogle B (1995) Effect of replacing whole boiled soya beans with azolla in the diets of growing ducks. *Livest. Res. Rural Dev.*, **7**: 1-11.
- Banerjee GC (2000) Poultry. Oxford and IBH Publishing, India.
- Basak B, Pramanik MAH, Rahman MS, Tarafdar SU and Roy BC (2002) Azolla (*Azolla pinnata*) as a feed ingredient in broiler ration. *Int. J. Poult. Sci.* **1**:29-34. DOI: 10.3923/ijps.2002.29.34
- Bhuyan MAH, Hasanat MR, Ali MA and Rahman MA (1998) Effect of feeding Azolla (*Azolla pinnata*) on the performance of broiler. *Bangladesh J. Anim. Sci.* **27**: 77-82.
- Cambel IM (1984) Growth Performance of Broilers Fed with Varying Levels of Azolla Meal. Kabacan, North Contabato, Philippines, pp: 66.
- Castilo LS, Gerpacio AL and Paseual FSD (1981) Exploratory Studies on Azolla and Fermented Rice Hulls in Broiler Diets. College, Leguna, Philippines, pp: 6.
- Fasuyi AO, Dairo FAS and Adeniji AO (2007) Protein Supplementary Quality of Tropical Vegetable (*Amaranthus cruentus*) Leaf Meal in Broiler Starter Diets: Bionutritional Evaluation. *International Journal of Agricultural Research*. **2**: 976-986. DOI: 10.3923/ijar.2007.976.986
- FTF (2010) Nepal FY 2010 Implementation Plan. Feed the Future. Retrieved on Feb. 20, 2012 from [http://www.feedthefuture.gov/sites/default/files/country/strategies/files/FTF\\_2010\\_Implementation\\_Plan\\_Nepal.pdf](http://www.feedthefuture.gov/sites/default/files/country/strategies/files/FTF_2010_Implementation_Plan_Nepal.pdf)
- Hossain MB, Mian MH, Hashem MA, Islam MZ and Shamsuddoha ATM (2001) Use of Azolla as Biofertilizer for Cultivation of BR 26 Rice in Aus Season. *Journal of Biological Sciences*. **1**: 1120-1123. DOI: 10.3923/jbs.2001.1120.1123
- Johari DC and Hussain KQ (1996) Commercial Broiler Production. International Book Distributing Company, Lucknow, Pages: 69.
- Kamalasanana P, Premalatha S, Rajamony S (2002) Azolla: A Sustainable Feed Substitute for Livestock. Tamilnadu, India.
- Krishnakumar S, Saravanan A, Ramesh K, Natarajan SK, Veerabadran V and Mani S (2005) Organic Farming: Impact on Rice (*Oryza sativa* L.) Productivity and Soil Health. *Asian Journal of Plant Sciences* **4**: 510-512. DOI: 10.3923/ajps.2005.510.512
- Muzlar AJ, Slinger SJ and Burton JH (1978) Chemical composition of aquatic macrophytes III. Mineral composition of fresh water macrophytes and their potential for mineral nutrient removal from lake water. *Canadian J. Plant Sci.* **58**: 851-862. DOI: 10.4141/cjps78-124
- Olorunfemi TOS, Aderibigbe FM, Alese BK and Fasakin EA (2006) Utilization of Duckweed (*Lemna paucicostata*) in Least-cost Feed Formulation for Broiler Starter: A Linear Programming Analysis. *Information Technology Journal*, **5**: 166-171. DOI: 10.3923/itj.2006.166.171
- Querubin LJ, Alcantara PF, Luis ES and Princesa AO (1986) Chemical composition and feeding value of Azolla in broiler ration. *Philippines J. Vet. Anim. Sci.* **12**: 65-65.
- Samanta G and Tamang Y (1995) Feeding value of azolla (*Azolla pinnata*) in goats. *Ann. Zootech.* **44**: 62-62. DOI: 10.1051/animres:19950532
- Sheeno TP and Sahu NP (2006) Use of Freshwater Aquatic Plants as a Substitute of Fishmeal in the Diet of Labeo rohita Fry. *Journal of Fisheries and Aquatic Science*. **1**: 126-135. DOI: 10.3923/jfas.2006.126.135
- Tahmid AH, Malek MA, Ahmed MU, Sota T and Tetsushi Y (2002) Influence of Azolla pinnata in Combination with Mustard Oil Cake on the Galling of *Meloidogyne javanica* and the Growth of Eggplant. *Pakistan Journal of Biological Sciences*. **5**: 665-668. DOI: 10.3923/pjbs.2002.665.668
- USAID (2008) The Nepal Poultry Value Chain and Avian Influenza. Retrieved on Feb 20, 2012 from [http://avianflu.aed.org/docs/supply\\_chain/Nepal\\_ValueChain\\_AI\\_Jun2008.pdf](http://avianflu.aed.org/docs/supply_chain/Nepal_ValueChain_AI_Jun2008.pdf)