



# International Journal of Applied Sciences and Biotechnology

A Rapid Publishing Journal

**ISSN 2091-2609**

## Indexing and Abstracting

CrossRef, Google Scholar, Global Impact Factor, Genamics, Index Copernicus, Directory of Open Access Journals, WorldCat, Electronic Journals Library (EZB), Universitätsbibliothek Leipzig, Hamburg University, UTS (University of Technology, Sydney): Library, International Society of Universal Research in Sciences (EyeSource), Journal Seeker, WZB, Socolar, BioRes, Indian Science, Jadoun Science, Jour-Informatics, Journal Directory, JournalTOCs, Academic Journals Database, Journal Quality Evaluation Report, PDOAJ, Science Central, Journal Impact Factor, NewJour, Open Science Directory, Directory of Research Journals Indexing, Open Access Library, International Impact Factor Services, SciSeek, Cabell's Directories, Scientific Indexing Services, CiteFactor, UniSA Library, InfoBase Index, Infomine, Getinfo, Open Academic Journals Index, HINARI, etc.

**CODEN (Chemical Abstract Services, USA): IJASKD**

Vol-3(4) December, 2015

**Available online at:**

<http://www.ijasbt.org>  
&  
<http://www.nepjol.info/index.php/IJASBT/index>



**Impact factor\*: 1.422**  
**Scientific Journal Impact factor#: 3.419**  
**Index Copernicus Value: 6.02**  
**IBI Factor 2015\*\*: 4.19**

\*Impact factor is issued by Universal Impact Factor. Kindly note that this is not the IF of Journal Citation Report (JCR).  
#Impact factor is issued by SJIF INNO SPACE; \*\*Impact factor is issued by INFOBASE INDEX.

For any type of query and/or feedback don't hesitate to email us at: [editor.ijasbt@gmail.com](mailto:editor.ijasbt@gmail.com)



Research Article

## A ROLE OF CALCIUM SIGNALING GENES IN HETEROKARYON INCOMPATIBILITY IN *NEUROSPORA CRASSA*

Ravi Gedela<sup>1\*</sup> and Ranjan Tamuli<sup>2</sup>

<sup>1</sup>Department of Bio-Science, Rajiv Gandhi University of Knowledge Technologies (RGUKT) Nuzvid-521201, India.

<sup>2</sup>Department of Biosciences and Bioengineering, Indian Institute of Technology, Guwahati-781039, India.

\*Corresponding author's email: ravibio531@rgukt.in

### Abstract

We have studied the Ca<sup>2+</sup>-signaling knockout mutants for their role in mating-type-associated heterokaryon incompatibility in *Neurospora crassa*. The found results showed on heterokaryons homokaryosis for ΔNCU05225, ΔNCU06366, ΔNCU06650, ΔNCU07075, and ΔNCU07966 Ca<sup>2+</sup>-signaling knockout mutants (*Neurospora crassa* unit number, NCU) displayed heterokaryon *het* compatibility; however heterokaryons heterokaryosis for ΔNCU05225, ΔNCU063665, ΔNCU06650, ΔNCU07075, and ΔNCU07966 mutants displayed *het* incompatibility like the wild-type control. In addition to that Two Ca<sup>2+</sup>-signaling knockout mutants ΔNCU02283, and ΔNCU09655 were tested for mating-type-associated heterokaryon incompatibility; these results showed, heterokaryons homokaryosis and heterokaryons heterokaryosis for ΔNCU02283, ΔNCU09655 mutants displayed *het* incompatibility. Cell death and hyphal compartmentation due to mating type associated incompatibility were confirmed by uptake of vital dye Evan's blue. Thus, these results of NCU05225, NCU06366, NCU06650, NCU07075, and NCU07966 Ca<sup>2+</sup>-signaling gene products could play a role in mating-type-associated heterokaryon incompatibility in *N. crassa*. In this article, we are reporting initially screened Ca<sup>2+</sup>-signaling gene deletion mutants of these five acts as recessive suppressors of mating type associated vegetative incompatibility in *N. crassa*.


**Keywords:** Ca<sup>2+</sup>-signaling genes; *Neurospora crassa* unit number (NCU); heterokaryon incompatibility; *Neurospora crassa*; non-self-recognition; vegetative incompatibility.

### Introduction

A cell containing different nuclear types in the same cytoplasm is called heterokaryon. Heterokaryon incompatibility refers to a condition when two nuclei of different genotypes cannot co-exist together in the same cytoplasm (Fig.1) (Staben and Yanofsky, 1990; Glass *et al.*, 1988, Debets and Griffiths, 1998; Worrall, 1997, Caten, 1972; Debets *et al.*, 1994; Van Diepeningen *et al.*, 1998). In *N. crassa*, both sexual and vegetative heterokaryon systems are present, sexual recognition is restricted by the mating type locus, whereas vegetative *het* incompatibility is genetically control by particular loci termed as *het* (for heterokaryon incompatibility) or *vic* (vegetative heterokaryon incompatibility)(Glass *et al.*, 1988; Ferreira, *et al.*,1998). When two fungal individuals of distinct *het* alleles come together, the consequential heterokaryotic cells are quickly damaged or strictly inhibited in their growth (Glass, 2006 and Marek *et al.*, 2003; Saupe, 2000) (Fig.1). This is analogous to histocompatibility system in invertebrates and major histocompatibility (MHC) in mammals (Xiang, 2002; Glass and Kaneko, 2003; Xiang, 2004). *N. crassa* has two mating (*mat*) types *mat A* and *mat a*. The coexistence of *mat A* and *mat a* during the vegetative

phase is lethal; however, coexistence of both mating types is necessary for the sexual development (Saupe, 2000; Garnjobst *et al.*, 1956, Wu, 2001; Sarkar *et al.*, 2002). These Two mating (*mat*) types *mat A* and *mat a*, alleles are differ in sequences although they occupy the same loci in different strains, and termed as 'idiomorph' (Glass *et al.*, 1988). The *A* idiomorph is 5301 base pairs that consists of *mat A-1*, *mat A-2* and *mat A-3* genes; in contrast, *a* idiomorph is 3235 base pair and contains only one ORF, *mat a-1*. The total number of *het* loci in *N. crassa* is 11, five of them the mating locus, *het-c*, *-d*, *-e*, and *i*, they have been recognized using enforced heterokaryons between nearly isogenic strains. Among them, three have been characterized at the molecular level, i.e. the mating type *mat* (*mat A* and *mat a*) locus, *het-c* and *het-6* (Glass *et al.*, 1988; Xiang, 2002; Glass and Kaneko, 2003; Xiang, 2004; Saupe, 2000; Wu, 2001; Sarkar *et al.*, 2002).

A non-self-recognition process called heterokaryon incompatibility (*het*) that operates during the vegetative and sexual phases of the filamentous in *N. crassa* (Glass *et al.*, 1988; Ferreira *et al.*, 1998; Garnjobst, *et al.*, 1956).



**Fig. 1:** Mechanism of heterokaryon incompatibility. Adapted from Gale Wichmann *et al.*, 2008.

### Heterokaryon Incompatibility in Other Filamentous Fungi

The fungal mating types which are tremendously dissimilar from each other, and do not show homology between strains of the opposite sex (as opposed to the allelic relationship in most polymorphic systems). The variance between dissimilar allelic forms of a *het* gene is generally widespread, but single-amino-acid differences can be sufficient to trigger incompatibility (Newmeyer, 1970; Glass *et al.*, 1988; Deleu, 1993). The *mat A-1* gene, encode a polypeptide of 288 amino acids region containing amino acids 90-104 has significant similarity to the MAT  $\alpha 1$  polypeptide of *Saccharomyces cerevisiae* (Glass *et al.*, 1990), this is required for the expression of the *het* incompatibility and sexual functions. The *mat a-1* encodes a 382 amino acids residues MAT a-1 polypeptide consisting of an HMG domain, and a separate region from amino acid residues 216-220 that confers vegetative incompatibility (*vic*) in *N. crassa* (Staben and Yanofsky, 1990; Glass *et al.*, 1990; Philley and Staben, 1994). In *Podospora anserine* a having two *het* loci, *het-s* and *het-e* are functional similar to the that of *het-c* and *het-6* of *N. crassa*, and *vib-1* (vegetative incompatibility block-1) of *Aspergillus nidulans*; but all of these genes are not homologous to each other (Ferreira *et al.*, 1998; Saupe, 2000; Sarkar *et al.*, 2002). The distinguish mechanism of *het* incompatibility reconcile by allele of differences at the *het-c* loci of *N. crassa* that inhibits phenotypic aspects of *het-c* vegetative incompatibility (Xiang, 2002; Glass and Kaneko, 2003; Xiang, 2004). The molecular description of *het* loci and *het* genes participating in the incompatibility effect has been achieved for two ascomycete's *N. crassa* and *P. anserine* (Smith *et al.*, 2000; Dementhon *et al.*, 2003; Xiang, 2002; Xiang, 2004; Glass and Kaneko, 2003; Saupe, 2000).

### Biological Significance of Heterokaryon Incompatibility

The biological significance of heterokaryon incompatibility, two different views was explained by subject. First, it has been proposed that heterokaryon incompatibility genes continue living limit nutrient situation, and heterokaryon development between different individuals (Debets, 1994; Glass *et al.*, 2004; Xiang, 2004). Second, the heterokaryotic cells strength boundary to the horizontal gene transfer of cytoplasmic infectious fundamentals elements such as senescence of plasmids, mycoviruses, transposes (Anagnostakis, 1977, 1983; Baidyaroy *et al.*, 2000; Biella *et al.*, 2002; Caten, 1972; Debets *et al.*, 1994; Van Diepeningen *et al.*, 1998; Hartl *et al.*, 1975; Hickey *et al.*, 2002; Glass *et al.*, 2004). On the other hand it has been proposed that mating type associated heterokaryon incompatibility might be maintaining out breeding by preventing the construction of different mating types of heterokaryons between siblings of the identical crosses (Debets and Griffiths, 1998; Worrall, 1997).

### Present Understanding of Mating Type-Associated Het Incompatibility in *N. Crassa*

*N. crassa* has two mating (*mat*) types, *mat A* and *mat a*, both mating types are essential for sexual development, however, coexistence during the vegetative phase is lethal, and therefore display incompatibility. The importance of mating type associated heterokaryon incompatibility, as anticipated for other *het* genes, the appearance of mating type allied incompatibility might be chance (Sarkar *et al.*, 2002). The *tol* gene (*tol* gene, for tolerant) is a suppressor of mating-type-associated heterokaryon incompatibility in *N. crassa* (Saupe, 2000). The  $Ca^{2+}$ -signaling genes regulates numerous processes secretion, sporulation, cytoskeletal organization, circadian rhythm, hyphal tip growth and hyphal branching in *N. crassa* (Fig. 1). Does Calcium signaling genes impact on mating-type-associated heterokaryon incompatibility in *N. crassa*? On this foundation, we designed the experiment by using auxotrophic marker for testing for complementation answer on  $Ca^{2+}$ -signaling genes in *N. crassa*; and we crossed with  $Ca^{2+}$ -signaling knockout mutants with auxotrophic marker *leu-3*, and *his-3* strains (Garnjobst, 1953; Adams *et al.*, 1987; Coenen, 1994).

## Materials and Methods

### Strains and Growth Conditions

Growth and maintenance of *Neurospora crassa* strains on Vogel's medium supplement with glucose were essentially as described in Davis and De Serres (1970). The  $Ca^{2+}$ -signaling knockout mutants, genotypes indicate as a *Neurospora crassa* unit number (NCU) (Table 1, and Table 2).

**Table 1:** Ca<sup>2+</sup>-signaling genes encodes proteins in *N. crassa*

S.N.	Ca <sup>2+</sup> -signaling genes	No. of amino acid	Encodes Name of Proteins
1	NCU02283	467	calcium/calmodulin-dependent protein kinase type I
2	NCU05225	674	mitochondrial NADH dehydrogenase
3	NCU06366	505	Ca <sup>2+</sup> /H <sup>+</sup> antiporter
4	NCU06650	186	Ca <sup>2+</sup> and /or CaM binding protein, a secretory phospholipase A2
5	NCU07075	508	CAX, Ca <sup>2+</sup> /H <sup>+</sup> exchanger
6	NCU07966	1110	Calcium transporting ATPase 3, a cation - ATPase
7	NCU09655	598 625	plasma membrane zinc ion transporter, phosphatidylinositol-4,5-bisphosphate phosphodiesterase gamma 2

**Note:** Ca<sup>2+</sup>-signaling genes encode proteins in *N. crassa*, available in site (<http://www.broadinstitute.org/annotation/genome/neurospora/>)

**Table 2:** list of strains used in this study.

S. No.	FGSC	Genotype	Source
1	12448 <i>a</i>	ΔNCU02283 <i>a</i>	FGSC
2	12449A	ΔNCU02283 A	FGSC
3	11405 <i>a</i>	ΔNCU05225 <i>a</i>	FGSC
4	11406A	ΔNCU05225A	FGSC
5	11407 <i>a</i>	ΔNCU06366 <i>a</i>	FGSC
6	114084	ΔNCU06366 A	FGSC
7	11246 <i>a</i>	ΔNCU06650 <i>a</i>	FGSC
8	11247A	ΔNCU06650A	FGSC
9	11248 <i>a</i>	ΔNCU07075 <i>a</i>	FGSC
10	11249 A	ΔNCU07075A	FGSC
11	11409 A	ΔNCU07966A	FGSC
12	11410 A	ΔNCU07966A	FGSC
13	11271 <i>a</i>	ΔNCU09655 <i>a</i>	FGSC
14	11272A	ΔNCU09655A	FGSC
15	1321 A	<i>leu-3A</i>	FGSC
16	3740 <i>a</i>	<i>leu-3a</i>	FGSC
17	6103 A	<i>his-3A</i>	FGSC
18	9716 <i>a</i>	<i>his-3a</i>	FGSC

**Table 3:** List of Ca<sup>2+</sup>-signaling knockout mutants obtained intensity band by Southern hybridization

S. No	Gene	R. E	Pro be	Wild type bands	Knockout bands
1	NCU02283A	<i>Nc</i> <i>ol</i>	5F - 5R	10kb,9.5 kb	4kb
2	NCU02283 <i>a</i>	<i>Nc</i> <i>ol</i>	5F - 5R	10kb,9.5 kb	4kb
3	NCU05225A	<i>Sal</i> <i>I</i>	5F - 5R	0.8kb	5.5kb
4	NCU05225 <i>a</i>	<i>Sal</i> <i>I</i>	5F - 5R	0.8kb	5.5kb
5	NCU06366A	<i>Sa</i> <i>c I</i>	5F - hph R	5kb,1.5k b	6kb
6	NCU06366 <i>a</i>	<i>Sa</i> <i>c I</i>	5F - hph R	5kb,4kb	6kb
7	NCU06650A	<i>Sa</i> <i>c I</i>	5F - hph R	6kb,0.8k b	6.5kb,3kb, and 0.8kb
8	NCU07075A	<i>Hi</i> <i>nd III</i>	3F - 3R	1kb	5kb,1kb
9	NCU07075 <i>a</i>	<i>Hi</i> <i>nd III</i>	3F - 3R	1kb	5kb,1kb
10	NCU07966A	<i>Hi</i> <i>nd III</i>	5F - 5R	9.5kb,8k b,and 6kb	10kb,8kb, and 0.7kb
11	NCU07966 <i>a</i>	<i>Hi</i> <i>nd III</i>	5F - 5R	10kb,8kb ,and 5kb	-
12	NCU09655A	<i>Xm</i> <i>a I</i>	5F - 5R	10kb,6kb ,and 5.5kb	5.5kb,4kb,2 kb,and 0.7kb
13	NCU09655 <i>a</i>	<i>Xm</i> <i>a I</i>	5F - 5R	10kb,6kb ,and 5.5kb	5.5kb,4kb,2 kb,and 0.7kb

The strains were obtained from the Fungal Genetic Stock Centre (FGSC), Kansas City, Missouri, USA. In this study we used two auxotrophic marker strains *leu-3* and *his-3* (*leu-3*, which is deficient to isopropylmalate synthetase, isopropylmalate dehydrogenase and isopropylmalate isomerase; *his-3*, histidinol dehydrogenase, phosphoribosyl-ATP-pyrophosphohydrolase and phosphoribosyl-AMP-cyclohydrolase) respectively; two auxotrophic strains *leu-3* and *his-3* were supplemented in the media with leucine (mg/ml) and histidine(mg/ml) nutrition for their proper growth. The crosses were performed by on Synthetic Crossing media 1X (SCM), and Sorbose-glucose-fructose media 1X (FGS) were used for germinating progeny/ascospore. The antibiotic hygromycin used at a working concentration 220 μg/ml for selecting Ca<sup>2+</sup>-signaling knockout mutants strains from their colonies.

**Test for Heterokaryon Incompatibility**

Strains with auxotrophic markers were cultured in Vogel's medium with required supplements. Conidial suspension of two auxotrophic strains, shown in below, were mixed and placed on a Petridis containing Vogel's glucose media without any supplement.

HO [*leu-3*; ΔNCU02283A) + (*his-3*; ΔNCU02283a), C1 [*leu-3*; NCU02283A) + (*his-3*;ΔNCU02283a], C2 [*leu-3*; ΔNCU02283A) + (*his-3*; NCU02283a], and C3 [*leu-3*; NCU02283A) + (*his-3*; NCU02283a] i.e. Ho: Heterokaryon homokaryosis and Heterokaryon heterokaryosis knockout test controls and well wild type control, similarly we examined all genes. That is, HO [(*ko1+*;mat A) + (*ko1+*;mat a)] are viable, whereas test controls C1[(*ko1+*; mat A) + (*ko1-*;mat a)], C2 [(*ko1-*; mat A) + (*ko1+*; mat a)]; and C3 [(*ko1-*; mat A) + (*ko1-*; mat a)] as a wild type are inviable (Table4, Fig. 2, Fig. 4a).

**Microscopic Analysis with Evan's Blue Staining**

Sterile pieces of cellophane were spread on top of the surface of solid Vogel's glucose agar medium. Heterokaryons were enforced by co-inoculating conidia of two strains grown on the cellophane for 2 days. The cellophane contain hyphae was peeled off from the surface of the medium and stained with 1% Evan's Blue dye (GAFF, 1971; JACOBSON *et al.*, 1998). The hyphae were stained for 15 to 25 minutes, and after that the cellophane was placed in a Buchner funnel containing a piece of filter paper pre-wetted with sterile water. Sterile water was then gently pipette over the cellophane and very weak vacuum was applied in order to clean off overload dye. The cellophane was then sited on a glass slide, with 5% glycerol to avoid drying, and observed under the bright field phase contrast microscope (Jacobson *et al.*, 1998; Wu and Glass 2001; Xiang, 2002; Glass, 2002).


**Table 4:** Test for heterokaryon incompatibility activity

S. No.	Heterokaryon type	<i>het</i> incompatibility
1	[( <i>leu-3</i> ;ΔNCU02283A(105))+( <i>his-3</i> ;ΔNCU02283a(86))]	No
2	[( <i>leu-3</i> ;NCU02283A(104))+( <i>his-3</i> ;ΔNCU02283a(86))]	Yes
3	[( <i>leu-3</i> ;ΔNCU02283A(105))+( <i>his-3</i> ;NCU02283 a(85))]	Yes
4	[( <i>leu-3</i> ;NCU02283A(104))+( <i>his-3</i> ;NCU02283a(85))]	Yes
5	[( <i>leu-3</i> ;ΔNCU05225a(93))+( <i>his-3</i> ;ΔNCU05225A(63))]	No
6	[( <i>leu-3</i> ;NCU05225a(89))+( <i>his-3</i> ;ΔNCU05225 A(63))]	Yes


S. No.	Heterokaryon type	<i>het</i> incompatibility
7	[( <i>leu-3</i> ;ANCU05225a(93))+( <i>his-3</i> ;NCU05225 A(64))]	Yes
8	[( <i>leu-3</i> ;NCU05225a(89))+( <i>his-3</i> ;NCU05225 A(64))]	Yes
9	[( <i>leu-3</i> ;ANCU06366A(13))+( <i>his-3</i> ;ΔNCU06366 a(6))]	No
10	[( <i>leu-3</i> ;ANCU06366A(13))+( <i>his-3</i> ;NCU06366 a(38))]	Yes
11	[( <i>leu-3</i> ;NCU06366A(14))+( <i>his-3</i> ;ΔNCU06366 a(6))]	Yes
12	[( <i>leu-3</i> ;NCU06366A(14))+( <i>his-3</i> ;NCU06366 a(38))]	Yes
13	[( <i>leu-3</i> ;ANCU06650a(37))+( <i>his-3</i> ;ΔNCU06650A(17))]	No
14	[( <i>leu-3</i> ;ANCU06650a(65))+( <i>his-3</i> ;NCU06650 A(41))]	Yes
15	[( <i>leu-3</i> ;NCU06650a(66))+( <i>his-3</i> ; ΔNCU06650 A(44))]	Yes
16	[( <i>leu-3</i> ;NCU06650a(66))+( <i>his-3</i> ;NCU06650 A(41))]	Yes
17	[( <i>leu-3</i> ;ANCU07075 a(22))+( <i>his3</i> ;ΔNCU07075A(19))]	No
18	[( <i>leu-3</i> ;ΔNCU07075a(22))+( <i>his-3</i> ;NCU07075A(24))]	Yes
19	[( <i>leu-3</i> ;NCU07075a(25))+ ( <i>his-3</i> ;ΔNCU07075A(19))]	Yes
20	[( <i>leu-3</i> ;NCU07075a(25))+( <i>his-3</i> ;NCU07075A(24))]	Yes
21	[( <i>leu-3</i> ;ANCU07966a(44))+( <i>his-3</i> ;ΔNCU07966A(34))]	No
22	[( <i>leu-3</i> ;NCU07966a(31))+( <i>his-3</i> ;ΔNCU07966A(34))]	Yes
23	[( <i>leu-3</i> ; ΔNCU07966a(44))+( <i>his-3</i> ;NCU07966A(8))]	Yes
24	[( <i>leu-3</i> ;NCU07966a(31))+( <i>his-3</i> ;NCU07966 A(8))]	Yes
25	[( <i>leu-3</i> ;ANCU09655a(97))+( <i>his-3</i> ;ΔNCU09655A(70))]	Yes
26	[( <i>leu-3</i> ;NCU09655a(96))+( <i>his-3</i> ; ΔNCU09655A(70))]	Yes
27	[( <i>leu-3</i> ; ΔNCU09655a(97))+( <i>his-3</i> ;NCU09655A(78))]	Yes
28	[( <i>leu-3</i> ;NCU09655a(96))+( <i>his-3</i> ;NCU09655A(78))]	Yes

## Results and Discussion


In primarily, we screened for heterokaryon incompatibility, heterokaryons homokaryosis and heterokaryons heterokaryosis of 20  $Ca^{2+}$ -signaling knockout mutants out of 48 merely in Yeast, Peptone and D-glucose (YPD) media, results showed *het* incompatibility. On this foundation, we crossed  $\Delta$ NCU05225,  $\Delta$ NCU06366,  $\Delta$ NCU06650,  $\Delta$ NCU07075, and  $\Delta$ NCU07966  $Ca^{2+}$ -signaling knockout mutants with the opposite mating types of *leu-3* and *his-3* auxotrophic marker strains (Table1 and Table 2). We found that the result heterokaryons homokaryosis for  $\Delta$ NCU05225,  $\Delta$ NCU06366,  $\Delta$ NCU06650,  $\Delta$ NCU07075, and  $\Delta$ NCU07966 mutants overcomes *het* incompatibility, therefore display mating type associated *het* compatibility and form vigours conidia on Petridis (Fig.2, Test: HO, Table4). Whereas, heterokaryons heterokaryosis for  $\Delta$ NCU05225,  $\Delta$ NCU06366,  $\Delta$ NCU06650,  $\Delta$ NCU07075, and  $\Delta$ NCU07966 mutants continue to showed mating type associated *het* incompatibility and showed killing of conidia on Petridis (Fig. 2, Test: HO, Test controls: C1 and C2; wild type control: C3; Table4). These results indicate that the knockout mutants of  $Ca^{2+}$ -signaling genes NCU05225, NCU06366, NCU06650, NCU07075, and NCU07966 suppressing the mating type associated *het* incompatibility in a recessive manner, and it showed genetically complementation in the heterokaryons of homokaryosis. The  $Ca^{2+}$ -signaling genes NCU05225, NCU06366, NCU06650, NCU07075, and NCU07966 encode proteins mitochondrial NADH dehydrogenase,  $Ca^{2+}/H^{+}$  antiporter,  $Ca^{2+}$  and /or CaM binding protein a secretory phospholipase A2, CAX  $Ca^{2+}/H^{+}$  exchanger, and Calcium transporting ATPase 3 respectively (Table1). We have also tested two additional  $Ca^{2+}$ -signaling knockout mutants  $\Delta$ NCU02283,  $\Delta$ NCU09655. We found the results in both the cases heterokaryons of homokaryosis and heterokaryons heterokaryosis for  $\Delta$ NCU02283,  $\Delta$ NCU09655 mutants display mating type associated *het* incompatibility like the wild-type control (Fig.3.3a, Test: HO, test controls: C1 and C2; wild type control: C3). The  $Ca^{2+}$ -signaling gene deletion mutants of these acts as recessive phenotypic expression of mating type associated *het* incompatibility. Therefore showed cell death on Petridis (Fig.3.3a, test: Ho, test controls: C1 and C2, wild type control: C3). This programmed cell death was confirmed by staining with vital dye Evan's Blue (Fig.3 and Fig. 4b, test: HO, test controls: C1 and C2, wild type control: C3). These  $Ca^{2+}$ -signaling genes NCU02283 and NCU09655 encodes protein calcium/calmodulin-dependent protein kinase type I and plasma membrane zinc ion transporter, phosphatidylinositol-4, 5-bisphosphate phosphodiesterase gamma 2 respectively (Table1).



**Fig. 2:**  $Ca^{2+}$ -signaling knockout mutant strains display mating type associated *het* compatibility activity in *N. crassa*. HO: [(*leu-3*;  $\Delta$ NCU06650 a (37)) + (*his-3*;  $\Delta$ NCU06650 A (17))]; C1: [(*leu-3*;  $\Delta$ NCU06650 a (65)) + (*his-3*; NCU06650 A (41))]; C2: [(*leu-3*; NCU06650 a (66)) + (*his-3*;  $\Delta$ NCU06650 A (44))]; C3: [(*leu-3*; NCU06650 a (66)) + (*his-3*; NCU06650 A (41))]. Here, heterokaryons homozygous for  $\Delta$ NCU05225,  $\Delta$ NCU06366,  $\Delta$ NCU06650,  $\Delta$ NCU07075, and  $\Delta$ NCU07966 mutants display *het* compatibility, and C1, C2 and C3 heterokaryons heterozygous for  $\Delta$ NCU05225,  $\Delta$ NCU06366,  $\Delta$ NCU06650,  $\Delta$ NCU07075, and  $\Delta$ NCU07966 mutants display *het* incompatibility.



**Fig. 3:** Microscopic analysis of cell death mediated by heterokaryon incompatible, using Evan's blue staining. HO: [(*leu-3*;  $\Delta$ NCU06650 a (37)) + (*his-3*;  $\Delta$ NCU06650 A (17))]; C1: [(*leu-3*;  $\Delta$ NCU06650 a (65)) + (*his-3*; NCU06650 A (41))]; C2: [(*leu-3*; NCU06650 a (66)) + (*his-3*;  $\Delta$ NCU06650 A (44))]; C3: [(*leu-3*; NCU06650 a (66)) + (*his-3*; NCU06650 A (41))]. Here, blue staining indicates killing due to mating type associated *het* incompatibility; dye retains the portion and remaining portion exclude. Heterokaryons homozygous for  $\Delta$ NCU05225,  $\Delta$ NCU06366,  $\Delta$ NCU06650,  $\Delta$ NCU07075, and  $\Delta$ NCU07966 mutants display *het* compatibility, therefore dye exclude within the portion.



**Fig. 4:** Two Ca<sup>2+</sup>-signaling genes NCU02283, NCU09655 do not suppress mating type associated het incompatibility

[a]. Mating type associated heterokaryon incompatibility was observed for Ho: [(leu-3; DNCU02283 a (105)) + (his-3; DNCU02283 A (86))]; C1: [(leu-3; NCU02283 a (104)) + (his-3; DNCU02283 A (86))]; C2: [(leu-3; DNCU02283 a (105)) + (his-3; NCU02283 A (85))]; C3: [(leu-3; NCU02283 a (104)) + (his-3; NCU02283 A (85))]; similarly for DNCU09655. Here, both the cases heterokaryons homozygous and heterozygous for □NCU02283, □NCU09655 mutants display het incompatibility. b). Microscopic analysis of heterokaryon incompatibility, using Evan's blue staining. Ho: [(leu-3; DNCU02283 a (105)) + (his-3; DNCU02283 A (86))]; C1: [(leu-3; NCU02283 a (104)) + (his-3; DNCU02283 A (86))]; C2: [(leu-3; DNCU02283 a (105)) + (his-3; NCU02283 A (85))]; C3: [(leu-3; NCU02283 a (104)) + (his-3; NCU02283 A (85))]; similarly for DNCU09655. Here, blue staining indicates killing due to mating type associated het incompatibility.]

The Ca<sup>2+</sup>-signaling knockout mutants were isolated from progeny by *hph* screening method, and we used strategy to test involvement of Ca<sup>2+</sup>-signaling genes in mating type of associate heterokaryons incompatibility. These Ca<sup>2+</sup>-signaling knockout mutants were confirmed by PCR (Fig. 5), using PCR primers (Table 5) and Southern hybridization methods (Fig. 5, 6). The Ca<sup>2+</sup>-signaling gene deletion mutants of these five acts as recessive suppressors of mating type associated *het* incompatibility; and result


reveals their role in mutant's recessive phenotypes of mating-type-associated heterokaryon incompatibility in *N. crassa*.

**Table 5:** List of Primers used in this study

S.NO	Gene	Primers
1	NCU02283	5F' AGG AGA AGT CTG AGA AGA GG 5R' CTA GAG GCA CCG TAC TAT CG
2	NCU05225	5F' TGT GAT TCA GGA TGT GGA GG 5R' GTT AGT GCA GCC AGT AAA GG
3	NCU06366	5F' CGG TAC ACT TGG TAA AGA GG 5R' AGT TGT AGA CAG GTA GGT GG
4	NCU06650	5F' TAC CTT ACC CAC CAG TAA CG 5R' CCT TCT CTT CTA TGT GCC AG
5	NCU07075	3F' GAG GTT TTC TGG TAG GGA GC 3R' GTA CCT CTA CCT AGC CTT GC
6	NCU07966	5F' AAG TTG AGT GTT CGT CCT CC 5R' GGT TCT TCT GTT CCT GTT CC
7	NCU09655	5F' CTG ACA GAG ATC TTG GAA GC 5R' GAT GAC TGA TGA CTG TGA CC



**Fig. 5:** PCR amplification of Ca<sup>2+</sup>-signaling genes to confirm the knockout mutants; 1) ΔNCU05225, 2) ΔNCU06366, 3) ΔNCU06650, 4) ΔNCU07075, 5) ΔNCU07966, 6) ΔNCU02283; and 7) ΔNCU09655. Here, 'M' is 1kb (NEB) marker.



**Fig. 6:**  $Ca^{2+}$ -signaling knockout mutant's confirmation by Southern hybridization. 1)  $\Delta$ NCU02283, 2)  $\Delta$ NCU05225, 3)  $\Delta$ NCU06366, 4)  $\Delta$ NCU06650, 5)  $\Delta$ NCU07075, 6)  $\Delta$ NCU07966 and 7)  $\Delta$ NCU09655 were digested with *NcoI*, *Sal I*, *Sac I*, *Sac I*, *Hind III*, *Hind III* and *Xma I* respectively, and resolved using of 1% agarose DNA was fragmented (Table 3). Here, 'M' is 1kb (NEB) marker; 1 (Wild types mating types *1m* and *1M*; mutant mating types *1a* and *1A*) 2, 3, 4, 5, 6 and 7 respectively. All probes obtained by gene specific PCR amplification with respective primers (except  $\Delta$ NCU06366,  $\Delta$ NCU06650 knockout probe, using primer *hph R*) 5F - 5R, 5F - 5R, 5F - *hphR*, 5F - *hphR*, 3F - 3R, 5F - 5R, and 5F - 5R (Table 4). Note that fragment resulting, respectively, from the digestion at the right and middle and the middle and left restriction sites of the intact gene is replaced in the disruption by the size of fragments duo to loss of the middle sites from the disrupted genes. Although the disruption bands is not seen as prominently in the DNA from wild type both mating types (*1m* and *1M*), it is seen in its hygromycin-resistant mutant both mating types (*1a* and *1A*).

## Discussion

The found results showed heterokaryons for homokaryosis  $\Delta$ NCU05225,  $\Delta$ NCU06366,  $\Delta$ NCU06650,  $\Delta$ NCU07075, and  $\Delta$ NCU07966 of suppression of mating type associated *het* compatibility (Fig. 3.1, HO, Table 4); and heterokaryons for heterokaryosis  $\Delta$ NCU05225,  $\Delta$ NCU06366,  $\Delta$ NCU06650,  $\Delta$ NCU07075, and  $\Delta$ NCU07966 mutants display *het* incompatibility like the wild-type control (Fig. 3.1, test: HO, test controls: C1 and C2, wild type control: C3; Table 4). That is, HO [(*ko1*;*mat A*) + (*ko1*;*mat a*)] are viable, whereas controls [(*ko1*+; *mat A*) + (*ko1*;*mat a*)] and [(*ko1*+; *mat A*) + (*ko1*+; *mat a*)] are inviable, likewise, for controls C1, C2, and C3 (Fig. 2, test: HO, test controls: C1 and C2, wild type control: C3; Table 4). Therefore  $\Delta$ NCU05225,  $\Delta$ NCU06366,  $\Delta$ NCU06650,  $\Delta$ NCU07075, and  $\Delta$ NCU07966  $Ca^{2+}$ -signaling genes have some significant role in *het* compatibility. An

additionally, same approach, we tested two more  $Ca^{2+}$ -signaling knockout mutants  $\Delta$ NCU02283,  $\Delta$ NCU09655 as control, for confirming of whether all calcium signaling genes showed heterokaryons compatibility (i.e. genetically complementation) or not. In this case two  $Ca^{2+}$ -signaling knockout mutants  $\Delta$ NCU02283,  $\Delta$ NCU09655 showed results heterokaryons for homokaryosis and heterokaryosis  $\Delta$ NCU02283,  $\Delta$ NCU09655 displayed *het* incompatibility (*vic*) as like the wild-type control (Fig. 4a, test: HO, test controls: C1 and C2, wild type control: C3). The microscopic analysis of the heterokaryons involving  $\Delta$ NCU05225,  $\Delta$ NCU06366,  $\Delta$ NCU06650,  $\Delta$ NCU07075, and  $\Delta$ NCU07966 mutants were confirmed by using the vital dye Evan's blue, as well  $\Delta$ NCU02283,  $\Delta$ NCU09655 knockout mutants. Which revealed neither hyphal compartmentation nor cell death showed for the heterokaryon homokaryosis (Fig. 3, test: HO), whereas in heterokaryon heterokaryosis showed cell death. During the

process of *het* incompatibility, some organelle modifications occurred like septal plugging, vacuolization, shrinkage of plasma membrane, organelle degradation, DNA fragmentation (Glass, 2006; Marek *et al.*, 2003) (Fig. 1), and accumulation of lipid bodies are common microscopic features associated with *het* incompatibility. Blue staining retaining was indication of killing due to mating type associated *het* incompatibility (Fig. 3 and 4b, test: HO, test controls: C1 and C2, wild type control: C3), and live cells portion exclude the dye (Fig. 3, test: HO). Both mating types are essential for sexual development, however, coexistence during the vegetative phase is lethal, and therefore display incompatibility (Fig. 1) (SAUPE, 2000; Garnjobst, *et al.*, 1956). The calcium signaling genes NCU05225 encode a 674aa residues of mitochondrial NADH dehydrogenase, NCU06366 encodes a 505aa residues of  $Ca^{2+}/H^{+}$  antiporter, NCU06650 encode a of 186aa residues of novel  $Ca^{2+}$  and/or CaM binding protein, a secretory phospholipase A2, NCU07075 encode a of 508 aa residues of CAX that is a  $Ca^{2+}/H^{+}$  exchange, and NCU07966 encodes a 1110aa residues of calcium transporting ATPase 3 (Table 1) respectively. In additionally other two  $Ca^{2+}$ -signaling genes encodes proteins like NCU02283 encode a 467aa residues of calcium/calmodulin-dependent protein kinase type I, and NCU09655 encodes plasma membrane zinc ion transporter (598aa) and phosphatidylinositol-4,5-bisphosphate phosphodiesterase gamma 2 (625aa) (Table 1) respectively. This calcium signaling genes encode protein might be involved in heterokaryons incompatibility process and further needs to be studied at the molecular level in *N. crassa*. Therefore, the result indicates suppression of mating type associated *het* incompatibility is not common phenomenon for all the  $Ca^{2+}$ -signaling knockout mutants in *N. crassa*.



## Conclusion

Previous reports suggest that the nuclei from two different genotypic strains are incompatible within the same cytoplasm (Garnjobst *et al.*, 1956; Saupe, 2000). In *N. crassa* at least 11 het loci exist. Five of them, the mating type locus and het-c, -d, -e and -I were originally identified using forced heterokaryons between nearly isogenic strains. Here, we reported five Ca<sup>2+</sup>-signaling knock mutant strains ΔNCU05225, ΔNCU06366, ΔNCU06650, ΔNCU07075, and ΔNCU07966 play important role in the mating-type-associated het incompatibility in *N. crassa* (Fig.2, test: HO, test controls: C1 and C2, wild type control: C3), additionally tested two more Ca<sup>2+</sup>-signaling knockout mutants ΔNCU02283, ΔNCU09655; in both the condition like heterokaryons homokaryosis and heterokaryosis of ΔNCU02283, ΔNCU09655 mutants display het incompatibility like the wild-type control (Fig. 4a, test: HO, test controls: C1 and C2, wild type control: C3). In *N. crassa* heterokaryon compatibility is shown by strains with identical genotypic class of progeny, whereas different genotypes of progeny are het incompatible. But the five Ca<sup>2+</sup>-signaling knockout mutant strains are het compatible suggesting that the knockout Ca<sup>2+</sup>-signaling gene may interact with the genes in het domain that result in induced mating type-associated het incompatibility in *N. crassa* (Fig.1.1 sup.info). Therefore, the mutants of

## References

- Aanen DK, Debets AJM, de Visser JAGM and Hoekstra RF (2008) The social evolution of somatic fusion. *BioEssays* **30**:1193-1203. DOI: 10.1002/bies.20840
- Adams G, Johnson N, Leslie JF and Hart LP (1987) Heterokaryons of *Gibberella zeae* formed following hyphal anastomosis or protoplast fusion. *Exp. Mycol.* **11**: 339-353. DOI: 10.1016/0147-5975(87)90022-3
- Altschul SF, Gish W, Miller W, Myers EW and Lipman DJ (1990) Basic local alignment search tool. *J. Mol. Biol.* **215**: 403-410. DOI: 10.1016/S0022-2836(05)80360-2
- Anagnostakis SL (1977) Vegetative incompatibility in *Endothia parasitica*. *Exp. Mycol.* **1**: 306-316. DOI: 10.1016/S0147-5975(77)80006-6
- Anagnostakis SL (1983) Conversion to curative morphology in *Endothia parasitica* and its restriction by vegetative compatibility. *Mycologia* **75**: 777-780. DOI: 10.2307/3792770
- Andrew P, Feinberg and Vogelstein B (1983) A technique for radiolabeling DNA restriction endonuclease fragments to high specific activity. *Anal Biochem.* **132**: 6-13. DOI: 10.1016/0003-2697(83)90418-9

heterokaryons homokaryosis identical *het* loci specificity displayed development rate, normal conidiation, and aerial hyphae formation, suggesting that the mutation was compatibility (Fig.2, HO) and heterokaryotic homokaryosis auxotrophic mutant strains growth in limited nutrient condition. Whereas heterokaryons heterokaryosis different *het* loci specificity displayed decrease in the development rate like lack of growth, normal conidiation, and aerial hyphae formation, suggesting that the mutation was incompatibility (Fig. 2, and Fig. 4a, test: HO, test controls: C1 and C2, wild type control: C3).

During the time of experiment setup, we have taken more care in mixing of conidial cells from other conidial strains contamination, and we verified specific knockout mutants mating type as well. We reported here the involvement of Ca<sup>2+</sup>-signaling gene in heterokaryon incompatibility as a phenotypic expression in *N. crassa* not yet reported and further needs to be studied at the molecular level.

## Acknowledgement

I thank my senior colleagues Ravi Kumar & Rekha for performing of Southern hybridization at CCMB, Hyderabad, India. I thank Fungal Genetics Stock Center (FGSC), University of Missouri, Kansas City, USA, for providing strains for free of charges; as well DBT, DST, MHRD and IITG for financial support. We are not paying attention on conflict of interest.

- Baidyaroy D, Glynn JM, Bertrand H (2000) Dynamics of asexual transmission of a mitochondrial plasmid in *Cryphonectria parasitica*. *Curr. Genet.* **37**: 257-267. DOI: 10.1007/s002940050527
- Balguerie A (2003) Domain organization and structure-function relationship of the HET-s prion protein of *Podospora anserina*. *EMBO J.* **22**: 2071-2081. DOI: 10.1093/emboj/cdg213
- Balguerie A (2004) The sequences appended to the amyloid core region of the HET-s prion protein determine higher-order aggregate organization in vivo. *J. Cell Sci.* **117**: 2599-2610. DOI: 10.1242/jcs.01116
- Barreau C, Iskandar M, Loubradou G, Levallois V and Begueret J (1998) The mod-A suppressor of nonallelic heterokaryon incompatibility in *Podospora anserina* encodes a proline-rich polypeptide involved in female organ formation. *Genetics* **149**: 915-926.
- Beadle GW and Coonradt VL (1944) Heterokaryosis in *Neurospora crassa*. *Genetics* **29**: 291-308.
- Biella S, Smith ML, Aist JR, Cortesi P, Milgroom MG (2002) Programmed cell death correlates with virus transmission in a filamentous fungus. *Proc. R. Soc. (London) - Ser. B: Biol. Sci* **269**: 2269-2276.

- Boye-Harnasch M and Cullin C (2006) A novel in vitro filter trap assay identifies tannic acid as an amyloid aggregation inducer for HET-s. *J. Biotechnol.* **125**: 222-230. DOI: 10.1016/j.jbiotec.2006.03.006
- Castro A, Lemos C, Falcao A, Glass NL and Videira A (2008) Increased resistance to complex I mutants to phytosphingosine-induced programmed cell death. *J. Biol. Chem.* **283**: 19314-19321. DOI: 10.1074/jbc.M802112200
- Caten CE (1972) Vegetative Incompatibility and Cytoplasmic Infection in Fungi. *J. Gen. Microbiol.* **72**: 221-229. DOI: 10.1099/00221287-72-2-221
- Coenen A, Debets AJM and Hoekstra RF (1994) Additive action of partial heterokaryon incompatibility (partial-het) genes in *Aspergillus nidulans*. *Curr. Genet.* **26**: 233-237. DOI: 10.1007/BF00309553
- Collins RA and Saville BJ (1990) Independent transfer of mitochondrial chromosomes and plasmids during unstable vegetative fusion in *Neurospora*. *Nature* **345**: 177-179. DOI: 10.1038/345177a0
- Coustou V (1999) Mutational analysis of the [Het-s] prion analog of *Podospora anserina*. A short N-terminal peptide allows prion propagation. *Genetics* **153**: 1629-1640
- Coustou V, Deleu C, Saupe S and Begueret J (1997) The protein product of the het-s heterokaryon incompatibility gene of the fungus *Podospora anserina* behaves as a prion analog. *Proc. Natl. Acad. Sci. USA* **94**: 9773-9778. DOI: 10.1073/pnas.94.18.9773
- Coustou-Linares V (2001) In vivo aggregation of the HET-s prion protein of the fungus *Podospora anserina*. *Mol. Microbiol.* **42**: 1325-35. DOI: 10.1046/j.1365-2958.2001.02707.x
- Dalstra HJ (2003) Sexual transmission of the [Het-s] prion leads to meiotic drive in *Podospora anserina*. *Proc. Natl. Acad. Sci. USA* **100**: 6616-6621. DOI: 10.1073/pnas.1030058100
- Dalstra HJ (2005) Non-mendelian inheritance of the HET-s prion or HET-s prion domains determines the het-S spore killing system in *Podospora anserina*. *Fungal Genet. Biol.* **42**: 836-47. DOI: 10.1016/j.fgb.2005.05.004
- Davis RH and De Serres FJ (1970) Genetic and microbiological research techniques for *Neurospora crassa*. *Methods Enzymol.* **17A**: 79-143. DOI: 10.1016/0076-6879(71)17168-6
- Debets AJM and Griffiths AJF (1998) Polymorphism of het-genes prevents resource plundering in *Neurospora crassa*. *Mycol. Res.* **102**: 1343-1349. DOI: 10.1017/S095375629800639X
- Debets F, Yang X and Griffiths AJ (1994) Vegetative incompatibility in *Neurospora*: its effect on horizontal transfer of mitochondrial plasmids and senescence in natural populations. *Curr. Genet.* **26**: 113-119. DOI: 10.1007/BF00313797
- Deleu C (1993) A single amino acid difference is sufficient to elicit vegetative incompatibility in the fungus *Podospora anserina*. *Genetics* **135**: 45-52.
- Deleu C, Turcq B and Begueret J (1990) Repa, a repetitive and dispersed DNA sequence of the filamentous fungus *Podospora anserina*. *Nucleic Acids Res* **18**: 4901-4903. DOI: 10.1093/nar/18.16.4901
- Dementhon K, Iyer G and Glass NL (2006) VIB-1 is required for expression of genes necessary for PCD in *Neurospora*. *Eukaryot Cell* **5**: 2161-2173. DOI: 10.1128/EC.00253-06
- Dementhon K, Paoletti M, Pinan-Lucarre B, Loubradou-Bourges N, Sabourin M and Saupe SJ (2003) Rapamycin mimics the incompatibility reaction in the fungus *Podospora anserina*. *Eukaryot. Cell* **2**: 2-246. DOI: 10.1128/EC.2.2.238-246.2003
- Felsenstein J (1981) Evolutionary trees from DNA sequences: a maximum likelihood approach. *J. Mol. Evol.* **17**: 368-376. DOI: 10.1007/BF01734359
- Ferreira ADV, An Z, Metzberg RL and Glass NL (1998) Characterization of mat A-2, mat A-3 and mat-A mating-type mutants of *Neurospora crassa*. *Genetics* **148**: 1069-1079.
- Ferreira ADV, Saupe S and Glass NL (1996) Transcriptional analysis of the mat A idiomorph of *Neurospora crassa* identifies two genes in addition to mat A-1. *Mol. Gen. Genet.* **250**: 767-774. DOI: 10.1007/BF02172989
- Fleissner A, Diamond S and Glass NL (2009) The *Saccharomyces cerevisiae* PRM1 homolog in *Neurospora crassa* is involved in vegetative and sexual cell fusion events, but also has post-fertilization functions. *Genetics* **181**: 497-510. DOI: 10.1534/genetics.108.096149
- Fleissner A, Leeder AC, Roca MG, Read ND and Glass NL (2009) Oscillatory recruitment of signaling proteins to cell tips promotes coordinated behavior during cell fusion. *Proc. Natl. Acad. Sci. USA* **106**: 19387-19392. DOI: 10.1073/pnas.0907039106
- Fleissner A, Sarkar S, Jacobson DJ, Roca MG, Read ND and Glass NL (2005) The so Locus Is Required for Vegetative Cell Fusion and Postfertilization Events

- in *Neurospora crassa*. *Eukaryot Cell* **4**: 920-430. DOI: 10.1128/EC.4.5.920-930.2005
- Fleissner A, Simonin AR and Glass NL (2008) Cell fusion in the filamentous fungus, *Neurospora crassa*. *Methods Mol. Biol.* **475**: 21-38. DOI: 10.1007/978-1-59745-250-2\_2
- Galagan JE (2003) The genome sequence of the filamentous fungus *Neurospora crassa*. *Nature* **422**: 859-68. DOI: 10.1038/nature01554
- Garnjobst L (1953) Genetic control of heterokaryosis in *Neurospora crassa*. *Am. J. Bot.* **40**: 607-614. DOI: 10.2307/2438448
- Garnjobst L (1955) Further analysis of genetic control of heterokaryosis in *Neurospora crassa*. *Am. J. Bot.* **42**: 444-448. DOI: 10.2307/2438792
- Garnjobst L and Wilson JF (1956) Heterokaryosis and protoplasmic incompatibility in *Neurospora crassa*. *Proc. Natl. Acad. Sci. USA* **42**: 613-618. DOI: 10.1073/pnas.42.9.613
- Glass NL and Kaneko I (2003) Fatal attraction: nonself recognition and heterokaryon incompatibility in filamentous fungi. *Eukaryot Cell* **1**: 1-8. DOI: 10.1128/EC.2.1.1-8.2003
- Glass NL and Kulda GA (1992) Mating-type and vegetative incompatibility in filamentous ascomycetes. *Annu. Rev. Phytopathol.* **30**: 201-224. DOI: 10.1146/annurev.py.30.090192.001221
- Glass NL and Lee L (1992) Isolation of *Neurospora crassa* Amating-type mutants by repeat-induced point (RIP) mutation. *Genetics* **132**: 125-133.
- Glass NL, Grotelueschen J and Metzberg RL (1990) *Neurospora crassa* Amating-type region. *Proc. Natl. Acad. Sci. USA* **87**: 4912-4916. DOI: 10.1073/pnas.87.13.4912
- Glass NL, Grotelueschen J and Metzberg RL (1990) *Neurospora crassa* A mating-type region. *Proc. Natl. Acad. Sci. USA* **87**: 4912-4916. DOI: 10.1073/pnas.87.13.4912
- Glass NL, Rasmussen C, Roca MG and Read ND (2004) Hyphal homing, fusion and mycelial interconnectedness. *Trends Microbiol.* **12**: 135-41. DOI: 10.1016/j.tim.2004.01.007
- Glass NL, Vollmer SJ, Staben C, Metzberg RL and Yanofsky C (1988) DNAs of the two mating type alleles of *Neurospora crassa* are highly dissimilar. *Science* **241**: 570-573. DOI: 10.1126/science.2840740
- Hartl DL, Dempster ER and Brown SW (1975) Adaptive significance of vegetative incompatibility in *Neurospora crassa*. *Genetics* **81**: 553-569.
- Hickey PC, Jacobson DJ, Read ND and Glass NL (2002) Live-cell imaging of vegetative hyphal fusion in *Neurospora crassa*. *Fungal Genet. Biol.* **37**: 109-119. DOI: 10.1016/S1087-1845(02)00035-X
- Hughes AL and Yeager M (1998) Natural selection at major histocompatibility complex loci of vertebrates. *Annu. Rev. Genet.* **32**: 415-435. DOI: 10.1146/annurev.genet.32.1.415
- Hutchison E, Brown S, Tian C and Glass NL (2009) Transcriptional profiling and functional analysis of heterokaryon incompatibility in *Neurospora crassa* reveals that reactive oxygen species, but not metacaspases, are associated with programmed cell death. *Microbiol* **155**: 3957-3970. DOI: 10.1099/mic.0.032284-0
- Jacobson DJ, Beurkensand K and Klomparens L (1998) Microscopic and ultrastructural examination of vegetative incompatibility in partial diploids heterozygous at the *het-s* locus in *Neurospora crassa*. *Fungal Genet. Biol.* **23**: 45-56. DOI: 10.1006/fgbi.1997.1020
- Jones DT, Taylor WR and Thornton JM (1992) The rapid generation of mutation data matrices from protein sequences. *Comp. Appl. Biosci.* **8**: 275-282. DOI: 10.1093/bioinformatics/8.3.275
- Koichiro T, Daniel P, Nicholas P, Glen S, Masatoshi N and Sudhir K (2011) Molecular Evolutionary Genetics Analysis Using Maximum Likelihood, Evolutionary Distance, and Maximum Parsimony Methods. *Mol. Biol. Evol.* **28**: 2731-2739. DOI: 10.1093/molbev/msr121
- Lew RR, Abbas Z, Anderca MI and Free SJ (2008) Phenotype of a mechanosensitive channel mutant, *mid-1*, in a filamentous fungus, *Neurospora crassa*. *Eukaryot Cell* **7**: 647-655. DOI: 10.1128/EC.00411-07
- Loubradou G, Bagueret J and Turcq B (1999) MOD-D, a Galpha subunit of the fungus *Podospora anserina*, is involved in both regulation of development and vegetative incompatibility. *Genetics* **152**: 519-28.
- Maddelein ML (2002) Amyloid aggregates of the HET-s prion protein are infectious. *Proc. Natl. Acad. Sci. USA* **99**: 7402-7407. DOI: 10.1073/pnas.072199199
- Malato L (2007) Role of Hsp104 in the propagation and inheritance of the [Het-s] prion. *Mol. Biol. Cell* **18**: 4803-4812. DOI: 10.1091/mbc.E07-07-0657

- Marchler B (2009) CDD: specific functional annotation with the Conserved Domain Database. *Nucleic Acids Res.* **37**: D205-D210. DOI: 10.1093/nar/gkn845
- Marek SM, Wu J, Glass NL, Gilchrist DG and Bostock RM (2003) Nuclear DNA degradation during heterokaryon incompatibility in *Neurospora crassa*. *Fungal Genet Biol.* **40**: 126-137. DOI: 10.1016/S1087-1845(03)00086-0
- Mathias S, Pena LA and Kolesnick RN (1998) Signal transduction of stress via ceramide. *Biochem. J.* **335**: 465-480. DOI: 10.1042/bj3350465
- Metzenberg RL and Glass NL (1990) Mating type and mating strategies in *Neurospora*. *Bio. Essays* **12**: 53-59. DOI: 10.1002/bies.950120202
- Micali OC and Smith ML (2005) A Nonself recognition gene complex in *Neurospora crassa*. *Genetics* **173**: 1991-2004. DOI: 10.1534/genetics.106.057562
- Mylyk OM (1975) Heterokaryon incompatibility genes in *Neurospora crassa* detected using duplication-producing chromosome rearrangements. *Genetics*. **80**: 107-124.
- Mylyk OM (1976) Heteromorphism for heterokaryon incompatibility genes in natural populations of *Neurospora crassa*. *Genetics* **83**: 275-84.
- Nazabal A (2003) Conformational transition occurring upon amyloid aggregation of the HET-s prion protein of *Podospora anserina* analyzed by hydrogen/deuterium exchange and mass spectrometry. *Biochemistry* **42**: 8852-8861. DOI: 10.1021/bi0344275
- Nazabal A (2005) Probing the structure of the infectious amyloid form of the prion-forming domain of HET-s using high resolution hydrogen/deuterium exchange monitored by mass spectrometry. *J. Biol. Chem.* **280**: 13220-13228. DOI: 10.1074/jbc.M413185200
- Nazabal A (2005a) High-resolution H/D exchange studies on the HET-s<sup>218-295</sup> prion protein. *J. Mass Spectrom* **40**: 580-590. DOI: 10.1002/jms.819
- Nazabal A and Schmitter JM (2006) Hydrogen-deuterium exchange analyzed by matrix-assisted laser desorption-ionization mass spectrometry and the HET-s prion model. *Methods Enzymol.* **413**: 167-181. DOI: 10.1016/S0076-6879(06)13009-8
- Newmeyer D (1970) A suppressor of the heterokaryon-incompatibility associated with mating type in *Neurospora crassa*. *Can. J. Genet. Cytol.* **12**: 914-926. DOI: 10.1139/g70-115
- Newmeyer D and Taylor CW (1967) A pericentric inversion in *Neurospora*, with unstable duplication progeny. *Genetics* **56**: 771-791.
- Perkins DD (1975) The use of duplication-generating rearrangements for studying heterokaryon incompatibility genes in *Neurospora*. *Genetics* **80**: 87-105.
- Perkins DD, Radford A, Newmeyer D and Bjorkman M (1982) Chromosomal Loci of *Neurospora crassa*. *Microbiol. Rev.* **46**: 426-570.
- Ren J, Wen L, Gao X, Jin C, Xue Y and Yao X (2009) DOG 1.0: illustrator of protein domain structures. *Cell Res.* **19**: 271-273. DOI: 10.1038/cr.2009.6
- Ritter C (2005) Correlation of structural elements and infectivity of the HET-s prion. *Nature.* **435**: 844-848. DOI: 10.1038/nature03793
- Sabate R (2007) Prion and non-prion amyloids of the HET-s prion forming domain. *J. Mol. Biol.* **370**: 768-783. DOI: 10.1016/j.jmb.2007.05.014
- Sabate R (2008) On the binding of Thioflavin-T to HET-s amyloid fibrils assembled at pH 2. *J. Struct. Biol.* **162**: 387-396. DOI: 10.1016/j.jsb.2008.02.002
- Sarkar S, Iyer G, Wu J and Glass NL (2002) Nonself recognition is mediated by HET-C heterocomplex formation during vegetative incompatibility. *EMBO J.* **18**: 4841-4850. DOI: 10.1093/emboj/cdf479
- Saupe S, Descamps C, Turcq B and Begueret J (1994) Inactivation of the *Podospora anserina* vegetative incompatibility locus *het-c*, whose product resembles a glycolipid transfer protein, drastically impairs ascospore production. *Proc. Natl. Acad. Sci. USA* **91**: 5927-5931. DOI: 10.1073/pnas.91.13.5927
- Saupe SJ, Kulda GA, Smith ML and Glass NL (1996) The product of the *het-C* heterokaryon incompatibility gene of *Neurospora crassa* has characteristics of a glycine-rich cell wall protein. *Genetics* **143**: 1589-1600.
- Shiu PKT and Glass NL (1999) Molecular characterization of *tol*, a mediator of mating-type-associated vegetative incompatibility in *Neurospora crassa*. *Genetics* **151**: 545-555.
- Siemer AB (2006) <sup>3</sup>C, <sup>15</sup>N resonance assignment of parts of the HET-s prion protein in its amyloid form. *J. Biomol. NMR* **34**: 75-87. DOI: 10.1007/s10858-005-5582-7
- Smith ML, Hubbard SP, Jacobson DJ, Micali OC and Glass NL (2000a) An osmotic-remedial, temperature-sensitive mutation in allosteric activity site of ribonucleotidoreductase in *Neurospora crassa*. *Mol.*

- Gen. Genet.* **262**: 1022-1035. DOI: 10.1007/PL00008645
- Smith ML, Micali OC, Hubbard SP, Mir-Rashed N, Jacobson DJ and Glass NL (2000) Vegetative incompatibility activity in the het-6 region of *Neurospora crassa* is mediated by two linked genes. *Genetics* **155**: 1095-1104.
- Staben C and Yanofsky C (1990) *Neurospora crassa* a mating-type region. *Proc. Natl. Acad. Sci. USA* **87**: 4917-4921. DOI: 10.1073/pnas.87.13.4917
- Sven SJ (2000) Molecular Genetics of Heterokaryon Incompatibility in Filamentous Ascomycetes. *Microbiol. Mol. Biol. Reviews* **64**: 489-502. DOI: 10.1128/MMBR.64.3.489-502.2000
- Taneja V (2007) A non-Q/N-rich prion domain of a foreign prion, [Het-s], can propagate as a prion in yeast. *Mol. Cell* **27**: 67-77. DOI: 10.1016/j.molcel.2007.05.027
- Thompson JD, Gibson TJ, Plewniak F, Jeanmougin F and Higgins DG (1997) The CLUSTAL\_X windows interface: flexible strategies for multiple sequence alignment aided by quality analysis tools. *Nucleic Acids Res.* **25**: 4876-4882. DOI: 10.1093/nar/25.24.4876
- Turcq B and Denayrolles M (1990) Isolation of two allelic incompatibility genes s and S of the fungus *Podospora anserina*. *Curr. Genet.* **17**: 297-303. DOI: 10.1007/BF00314876
- Van Diepeningen AD and Hoekstra RF (1997) Heterokaryon incompatibility blocks virus transfer among natural isolates of black Aspergilli. *Curr. Genet* **32**: 209-217. DOI: 10.1007/s002940050268
- Vellani TS, Griffiths AJF and Glass NL (1994) New mutations that suppress mating-type associated incompatibility in *Neurospora crassa*. *Genome* **37**: 249-255. DOI: 10.1139/g94-035
- Vogel HJ (1964) Distribution of lysine pathways among fungi: evolutionary implications. *Am. Nat.* **98**: 435-446. DOI: 10.1086/282338
- Wasmer C (2008) Amyloid fibrils of the HET-s (218-289) prion form a beta solenoid with a triangular hydrophobic core. *Science* **319**: 1523-1526. DOI: 10.1126/science.1151839
- Worrall JJ (1997) Somatic incompatibility in basidiomycetes. *Myc-ologia.* **89**: 24-36. DOI: 10.2307/3761169
- Wu J and Glass NL (2001) Identification of specificity determinants and the generation of alleles with novel specificity at the het-c heterokaryon incompatibility locus of *Neurospora crassa*. *Mol. Cell Biol.* **21**: 1045-1057. DOI: 10.1128/MCB.21.4.1045-1057.2001
- Xiang Q and Glass NL (2002) Identification of vib-1, a locus involved in vegetative incompatibility mediated by het-c in *Neurospora crassa*. *Genetics* **162**: 89-101.
- Xiang Q and Glass NL (2004) Chromosome rearrangements in isolates that escape from het-c heterokaryon incompatibility in *Neurospora crassa*. *Curr. Genet.* **44**: 329-323. DOI: 10.1007/s00294-003-0451-y
- Xiang Q and Glass NL (2004) The control of mating type heterokaryon incompatibility by vib-1, a locus involved in het-c heterokaryon incompatibility in *Neurospora crassa*. *Fungal Genet Biol.* **41**: 4-11076. DOI: 10.1016/j.fgb.2004.07.006
- Xiang Q, Rasmussen C and Glass NL (2002) The ham-2 locus, encoding a putative transmembrane protein, is involved in hyphal fusion in *Neurospora crassa*. *Genetics* **160**: 169-180.
- Yamazaki T (2008) Hydration effects on the HET-s prion and amyloid-beta fibrillous aggregates, studied with three-dimensional molecular theory of salvation. *Biophys. J.* **95**: 4540-4548. DOI: 10.1529/biophysj.107.123000