



ISSN: 2091-2749 (Print)
2091-2757 (Online)

Correspondence

Dr. Balakrishnan M Acharya
Department of Orthopedic and
Trauma Surgery, Patan Academy
of Health Sciences, Lalitpur,
Nepal
Email:
balakrishnanmacharya@pahs.
edu.np

Peer Reviewers

Asst. Prof. Dr. Ashis Shrestha
Patan Academy of Health
Sciences

Prof. Dr. Jay Narayan Shah
Patan Academy of Health
Sciences

Submitted

14 Oct 2018

Accepted

30 Nov 2018

How to cite this article

Balakrishnan M Acharya,
Pramod Devkota, Amrit
Shrestha, Abhishek Kumar
Thakur, Toya Raj Bhatta, Bidur
Gyawali. Outcome of surgical
management for the ruptured
Achilles tendon. Journal of
Patan Academy of Health
Sciences. 2018Dec;5(2):18-22.

Outcome of surgical management of ruptured Achilles tendon

Balakrishnan M Acharya,¹ Pramod Devkota,¹ Amrit Shrestha,² Abhishek Kumar Thakur,² Toya Raj Bhatta,³ Bidur Gyawali³

¹Associate Professor, ²Lecturer, ³Assistant Professor, Department of Orthopedic and Trauma Surgery, Patan Academy of Health Sciences, Lalitpur, Nepal

Abstract

Introductions: The incidence of Achilles tendon rupture has been increasing worldwide. The optimal management of this problem is controversial but the surgical intervention is being applied widely.

Methods: We retrospectively analyzed patients who had Achilles tendon (AT) rupture treated surgically at Patan Academy of Health Sciences from January 2010 to December 2015. Functional outcome assessment of ankle was done by using American Orthopedic foot and Ankle Society Score (AOFAS).

Results: Total 71 patients (male 45, female 26) had surgery for AT, 38 right and 33 left side. The mean age of the patient was 36.14 years (range 18-67 years). The mean AOFAS score was 83.32 (range 75-93) after first year follow up and 90.36 (range 80-99) after second year, difference was statistically significant ($p < 0.05$).

Conclusions: The study showed high AOFAS score for surgical management of ruptured Achilles tendon.

Key words: Achilles tendon rupture, American Orthopedic foot and Ankle Society Score (AOFAS),

Introductions

Achilles tendon (AT) is the strongest tendon in ankle and foot of the human body.¹ The AT injury may occur due to sports, accidental rupture, road traffic accidents, slipping of foot etc.²⁻³ Among all the tendon injuries that come in Orthopedic department, AT injury accounts for approximately 35%, mainly due to recreational activities, more in males of 30 to 40 years of age.⁴⁻⁶ It has a prolonged recovery and there is 10% to 30% reduction in the calf muscle strength and planter flexion.⁷⁻¹¹

The injury can be managed by both conservative and operative procedures. However, conservative management has shown poor outcome than surgical repair.¹⁰⁻¹² Surgical repair is increasingly used due to early recovery, lower re-rupture rate, better range of motion, better functional outcome and lower calf atrophy.¹²⁻¹³ This study aims to evaluate functional outcome of patient after surgical repair of AT.

Methods

This cross-sectional study was conducted at the department of orthopedics and trauma surgery of Patan Hospital, Patan Academy of Health Sciences (PAHS), Lalitpur, Nepal. The study was approved by the institutional review committee (IRC) of PAHS. The data of AT patients operated from January 2010 to December 2015 were analyzed. All patients above 16 years of age, who had AT rupture due to recreation activities were included in this study. Patient less than 15 years, diabetic foot, associated neurovascular injury and cut injury of AT by sharp object were excluded. Diagnosis was made mainly based on positive Thompson test¹⁴ and ultrasound of the AT. Radiological examinations were done to rule out associated fractures around the ankle. Operative procedure was performed by end-to-end repair of ruptured AT using two-stranded single Krackow suturing technique.¹⁵

Postoperatively, patients were put on below knee plaster cast for six weeks. Followup for functional outcome was done at one and two years. Since the patient do not come exactly at one and two years, their visit after one year and before two year were categorized as followup point A and the visit after two years as point B. Functional assessment of ankle was done by using American Orthopedic Foot and Ankle Society Score (AOFAS).¹⁶ Statistical analysis was carried out using the SPSS 20 software.

Results

There were 71 patients of AT rupture who were treated with surgery. Out of them male were 45 (63.3%) and female were 26 (36.7%). Rupture on right side was 38 (53.5%) and left side was 33 (46.5). The mean age of the patient was 36.14 years (18-67), mean duration of symptom presented was 6.08 days (1 to 18). The average hospital stay after surgery was 5.56 days (4 to 11), and mean follow up at point A was 14.14 months (12-18) and at point B was 30.76 months (24-39).

The mean AOFAS score was 83.32 (75-93) at point A and 90.36 (80-99) at point B. The difference was statistically significant ($p < 0.05$). No complain of difficulties in daily activities was reported. Fourteen patients (19.72%) complained that they were unable to perform running and jogging like they used to do before injury.

Four patients had swollen foot and cellulitis after surgery, 9/71 (12.68%) had infection at surgical sites which resolved on antibiotics and dressing. Ten patients complained of pain over scar site which resolved after physiotherapy. There were no other post-operative complications like re-rupture of AT, pain, numbness, neurovascular injury.



Figure 1. a) Achilles tendon rupture showing Thompson test positive in left leg, b) Intra operative picture showing Achilles tendon rupture with retracted plantaris tendon, c) Healed Scar (left leg)

Discussions

The mean AOFAS score was 83.32 (75-93) at point A, and 90.36 (80-99) at point B, the difference was statistically significant ($p < 0.05$). A similar study published in 2017¹⁷ got AOFAS score of 92.4 at one year; we were able to attain this score over two years period. This possibly suggest that the functional outcome of surgery is comparable but with longer rehabilitation that we need to look into. Our study also suggested statistically significant improvement between point of followup A and B. This suggest that longer rehabilitation, and comparison between these points was appropriate.

A prospective randomized trial reported the average AOFAS score of surgical treatment to be 86.7 and non-operative to be 55.0.¹⁸ The score for surgical treatment is consistent with our study, supporting that operative management is better than non-operative management as in our setup. Operative treatment of AT rupture provides superior results as compared to non-operative treatment in terms of complications like re-rupture, decreased plantar flexion strength of ankle muscle and its endurance.¹³ Our study shows 14 (19.7%) patients were unable to perform running and jogging like they used to do before injury. This is consistent with other studies which has reported that 85% to 100% patients returned to their recreational activities.¹⁹⁻²¹ This further supports surgical management is equally good in our setup like others.

In our study male patient got rupture of AT more often than female and more on right side. The other studies also showed the injury was more in male, however involved side varies in different studies.²²⁻²³ The surgical site infection rate in this study was 7/71 (12.67%), higher than others reporting 3% infection.²⁴ Possible reason could be the less hygienic environment and nutritional status of our patients. We did not encounter other complications like numbness of the foot and ankle, re-rupture of the repaired Achilles tendon. Numbness of the foot due to sural nerve injury in 9.6% and deep vein thrombosis (DVT) after surgery in 6.45% have been reported by some other studies.²²

The management of AT is a challenging task for Orthopedic surgeon.²⁵ After tendon rupture, the strength of plantar flexion of the ankle is reduced, and the patients are not able to perform a single-limb heel rise with the injured lower extremity.²⁶ This is regarded as the vital indication for surgical intervention in ruptured tendon. The return to daily activities after the surgery is an important indicator for the success of surgical intervention.²²

Conclusions

The overall outcome of the surgical repair of Achilles tendon rupture was safe and effective with good functional outcome and minimal minor complications.

References

1. Smith DW, Rubenson J, Lloyd D, et al. A conceptual framework for computational models of Achilles tendon homeostasis. *Wiley Interdiscip Rev Syst Biol Med*. 2013;5(5):523-38. DOI: 10.1002/wsbm.1229
2. Khan RJ, Fick D, Keogh A, Crawford J, Brammar T, Parker M. Treatment of acute Achilles tendon ruptures: a meta-analysis of randomized, controlled trials. *J Bone Joint Surg Am*. 2005;87(10):2202-10. [Web link](#)
3. Ingvar J, Tagil M, Eneroth M. Nonoperative treatment of Achilles tendon rupture: 196 consecutive patients with a 7% re-rupture rate. *Acta Orthop*. 2005;76(4):597-601. DOI: 10.1080/17453670510041619
4. Jozsa L, Kvist M, Balint BJ, et al. The role of recreational sport activity in Achilles tendon rupture: a clinical, pathoanatomical, and sociological study of 292 cases. *Am J Sports Med*. 1989;17(3):338-43. DOI: 10.1177/036354658901700305
5. Maffulli N, Ajis A, Longo UG, et al. Chronic rupture of tendo Achillis. *Foot Ankle Clin*. 2007;12(4):583-96. DOI: 10.1016/j.fcl.2007.07.007
6. Suchak AA, Bostick G, Reid D, et al. The incidence of Achilles tendon ruptures in Edmonton, Canada. *Foot Ankle Int*. 2005;26(11):932-6. DOI: 10.1177/107110070502601106
7. Heikkinen J, Lantto I, Flinkkila T, et al. Augmented compared with nonaugmented surgical repair after total Achilles rupture: results of a prospective randomized trial with thirteen or more years of follow-up. *J Bone Joint Surg Am*. 2016;98(2):85-92. DOI: 10.2106/JBJS.O.00496 PMID: 26791028
8. Olsson N, Nilsson-Helander K, Karlsson J, et al. Major functional deficits persist 2 years after acute Achilles tendon rupture. *Knee Surg Sports Traumatol Arthrosc*. 2011;19(8):1385-93. DOI: 10.1007/s00167-011-1511-3
9. Olsson N, Silbernagel KG, Eriksson BI, et al. Stable surgical repair with accelerated rehabilitation versus non-surgical treatment for acute Achilles tendon ruptures: a randomized controlled study. *Am J Sports Med*. 2013;41(12):2867-76. DOI: 10.1177/0363546513503282
10. Silbernagel KG, Steele R, Manal K. Deficits in heel-rise height and Achilles tendon elongation occur in patients recovering from an Achilles tendon rupture. *Am J Sports Med*. 2012;40(7):1564-71. DOI: 10.1177/0363546512447926
11. Willits K, Amendola A, Bryant D, et al. Operative versus non-operative treatment of acute Achilles tendon ruptures: a multicenter randomized trial using accelerated functional rehabilitation. *J Bone Joint Surg Am*. 2010;92(17):2767-75. DOI: 10.2106/JBJS.I.01401 PMID: 21037028
12. Kocher MS, Bishop J, Marshall R, et al. Operative versus nonoperative management of acute Achilles tendon rupture. *Am J Sports Med*. 2002;30(6):783-90. DOI: 10.1177/03635465020300060501
13. Strauss EJ, Ishak C, Jazrawi L, Sherman O, Rosen J. Operative treatment of acute Achilles tendon ruptures: an institutional review of clinical outcomes. *Injury*. 2007;38(7):832-8. DOI: 10.1016/j.injury.2006.06.005
14. Douglas J, Kelly M, Blachut P. Clarification of the Simmonds–Thompson test for rupture of an Achilles tendon. *Canadian Journal of Surgery*. 2009;52(3):E40-1. PMID: PMC2689757 [PDF](#)
15. Choi GW, Kim HJ, Lee TH, Park SH, Lee HS. Clinical comparison of the two-stranded single and four-stranded double Krackow techniques for acute Achilles tendon ruptures. *Knee Surg Sports Traumatol Arthrosc*. 2017;25(6):1878-83. DOI: 10.1007/s00167-016-4265-0.
16. Kitaoka HB, Alexander IJ, Adelaar RS, et al. Clinical rating systems for the ankle-hindfoot, midfoot, hallux, and lesser toes. *Foot Ankle Int*. 1994;15(7):349-53. DOI: 10.1177/107110079401500701
17. Li CG, Li B, Yang FY. Management of acute Achilles tendon rupture with tendon-bundle technique. *J Inte Med Reser*. 2017;45(1) 310-9. DOI: 10.1177/0300060516677928
18. Thordarson DB, Krieger LE. Operative vs. nonoperative treatment of intra-articular fractures of the calcaneus: a prospective randomized trial. *Foot Ankle Int*. 1996;17(1):2-9. DOI: 10.1177/107110079601700102
19. Mukundan C, El Hussein M, Rayan F, Salim J, Budgen A. “Mini-open” repair of acute tendon Achilles ruptures – the solution? *Foot Ankle Surg*. 2010;16(3):122-5. DOI: 10.1016/j.fas.2009.07.005
20. Garrido IM, Deval JC, Bosch MN, Mediavilla DH, Garcia VP, González MS. Treatment of acute Achilles tendon ruptures with Achillon device: clinical outcomes and kinetic gait

- analysis. *Foot Ankle Surg.* 2010;16(4):189-94. DOI: 10.1016/j.fas.2009.10.014
21. Suchak AA, Spooner C, Reid DC, Jomha NM. Postoperative rehabilitation protocols for Achilles tendon ruptures. a meta-analysis. *Clin Orthop Relat Res.* 2006;445:216-21. DOI: 10.1097/01.blo.0000203458.05135.74
22. Jallageas R, Bordes J, Daviet JC, Mabit C, Coste C. Evaluation of surgical treatment for ruptured Achilles tendon in 31 athletes. *Orthop Traumatol Surg Res.* 2013;99(5):577-84. DOI: 10.1016/j.otsr.2013.03.024
23. Lin Y, Yang L, Yin L, Duan X. Surgical strategy for the chronic Achilles tendon rupture. *Biomed Research Inter.* 2016. (article ID: 1416971) DOI: 10.1155/2016/1416971
24. Jildeh TR, Okoroha KR, Marshall NE, et al. Infection and rerupture after surgical repair of Achilles tendons. *Orthop J Sports Med.* 2018;6(5). DOI: 10.1177/2325967118774302
25. Cetti R, Christensen SE, Ejsted R, et al. Operative versus nonoperative treatment of Achilles tendon rupture: a prospective randomized study and review of the literature. *Am J Sports Med.* 1993;21(6):791-9. DOI: 10.1177/036354659302100606
26. Khiami F, Di Schino M, Sariali E, et al. Treatment of chronic Achilles tendon rupture by shortening suture and free sural triceps aponeurosis graft. *Orthop Traumatol Surg Res.* 2013;99(5):585-91. DOI: 10.1016/j.otsr.2013.03.021