

# Awareness regarding cardiac rehabilitation among patients with coronary heart disease attending a cardiac care centre, Kathmandu valley

Rosy Shrestha<sup>1</sup>, Sajeeb Shrestha<sup>2</sup>

<sup>1</sup> School of nursing, Chitwan medical college, Bharatpur, Nepal

<sup>2</sup> Department of public health, Central institute of science and technology, Kathmandu

**Corresponding Author:** Rosy Shrestha

School of nursing, Chitwan medical college, Bharatpur, Nepal

**Email address:** rosyshrestha2016@gmail.com

**Cite this article as:** Shrestha R, Shrestha S. Awareness regarding cardiac rehabilitation among patients with coronary heart disease attending a cardiac care centre, Kathmandu valley. Nepalese Heart Journal 2019; Vol 16(1), 47-50

**Received date:** 22nd March 2019

**Accepted date:** 16th April 2019

## Abstract

**Introduction:** Coronary heart disease (CHD) is gradually emerging as a leading cause of morbidity and mortality of many low middle income countries like Nepal. Cardiac rehabilitation awareness program has been proved to be effective for reducing the mortality as well as improving the quality of life among CHD patients. The aim of this study was to explore the awareness on cardiac rehabilitation (CR) in patients with CHD attending a cardiac care centre, Nepal.

**Method:** A descriptive cross sectional study design was used to examine 100 CHD patients attending out-patient departments of Shahid Gangalal National Heart Centre (SGNHC), Kathmandu, Nepal. Purposive sampling technique was used for data collection by face to face interview technique with self developed tool. Data was analyzed with descriptive and inferential statistics.

**Results:** Of all 100 respondents, 55.0% were male and the mean age was 53.23±14.22 years. The median score of awareness was 17 with interquartile range (IQR) 14.0-19.75 and majority (57.0%) of respondents were unaware about CR. The awareness regarding CR was found highest in awareness regarding CHD (75.0%) whereas lowest score was found in time and duration needed for exercise per week (14.0%). The significant influencing variables were age, education status, duration of treatment and participation in CR program for CHD patients.

**Conclusion:** The awareness on CR program in CHD patients in Nepal is not optimal, especially among 54 and above age group, illiterate people, those receiving treatment equal & more than 1 year duration and the CHD patients who didn't get chance to participate on awareness programs. Hence, it is strongly recommended that health professionals including nurses should organize and promote CR programs including counseling session to improve the awareness level and ultimately enhance quality of life of CHD patients.

**Key words:** Awareness, Cardiac Rehabilitation, Coronary heart disease

DOI: <https://doi.org/10.3126/njh.v16i1.23899>

## Introduction

Coronary heart disease (CHD) is the leading cause of morbidity & mortality worldwide, accounting for over one-quarter of all deaths in 2001 and in adult population deaths under 65 years (33%).<sup>1-2</sup> And it is forecasted that by 2020 CHD will be a major burden of disease worldwide.<sup>3</sup> A person with CHD presents on a continuum of events that includes angina, myocardial infarction (MI), and ischemic heart failure.<sup>4</sup> CHD has become a potential time bomb causing deaths in low and middle income countries like Nepal, where preventive measures have not been effective.<sup>5</sup> It is gradually emerging as one of the major health challenges and prevalence in Eastern region of Nepal was 6%.<sup>6</sup> Outpatient and inpatient rehabilitation are an essential part of tertiary prevention for the long-term success of medical treatment and it reduces both cardiovascular and total mortality rates for patients with CHD. It is also reported that 39.0% of CHD patients had clear indication for rehabilitation for their inpatient hospital stay.<sup>7</sup> Those who did not undergo

rehabilitation were more likely to develop diabetes mellitus, arterial hypertension, or peripheral arterial occlusive disease and had lower left ventricular ejection fractions than those who did undergo rehabilitation. Patients with CHD can benefit from cardiac rehabilitation (CR) programs.<sup>8</sup> Therefore this study was conducted with the purpose of investigating awareness on cardiac rehabilitation in CHD patients.

## Methods

A descriptive cross-sectional study design was used to examine 100 CHD patients attending outdoor department of Shahid Gangalal National Heart Center (SGNHC), Bansbari, Kathmandu, Nepal during November to December, 2015.. Non-probability purposive sampling technique was used to select sample, who were clinically diagnosed case of coronary heart disease (Myocardial Infarction, Angina Pectoris and Ischemic heart failure) after angiography for more than three months and subjects with 30 years and above age group were included whereas those who were unable and unwilling to participate

were excluded from this study. The pretested (Cronbach  $\alpha = 0.78$ ) tool was used for data collection which consisted three parts: demographic (total items-9) and disease related variables (total items-5) and awareness related cardiac rehabilitation (total items-17). Data was collected by face to face interview schedule at outdoor department of SGNHC, Bansbari, Kathmandu. Ethical clearance was taken from Institutional Review Board (IRB) of Institute of Medicine (IOM), Tribhuvan University, Kathmandu, Nepal. Informed consent (verbal and written as applicable) was obtained from each respondent prior to data collection. Data was analyzed using Statistical Package for the Social Sciences (SPSS) version 20.0. Descriptive (Percentage, frequency, mean and standard deviation, median, interquartile range (IQR) and inferential (Chi-square) statistic was used to find out association between level of awareness and selected variables. The level of awareness was classified into two categories as an aware and unaware based on median value (17) by Shapiro wilk test for normality. The median value of overall awareness was 17 (57.0%).

## Results

Table 1 shows that the majority of the respondents were aged 45-59 years (34.0%), male (55.0%), resided in urban (52.0%), were living with family (76.0%), Brahmin/Chhetri (59.0%), were Hindu (84.0%), belonged to joint family (51.0), literate (58.0), and housework as an occupation (35.0).

TABLE 1 - Socio-demographic Characteristics of the Respondents

n=100	
Variables	Frequency/Percent
<b>Age group (in years)</b>	
30-44	29.0
45-59	34.0
60-74	26.0
>74	11.0
Mean age $\pm$ SD= 53.23 $\pm$ 14.22; Min =30 & Max 88 years	
<b>Sex</b>	
Male	55.0
Female	45.0
<b>Place of Residence</b>	
Rural	48.0
Urban	52.0
<b>Living Status</b>	
Living with family	76.0
Living single*	24.0
<b>Ethnicity</b>	
Brahmin/Chhetri	59.0
Janajati	35.0
Dalit	6.0
<b>Religion</b>	
Hindu	84.0
Non-Hindu	16.0
<b>Type of Family</b>	
Nuclear	49.0
Joint	51.0

Education	
Literate	58.0
Illiterate	42.0
<b>Occupation*</b>	
Farming	21.0
Housework	35.0
Service	23.0
Business	21.0
* Included unmarried, divorced, widower/widow; **Included household activities like cooking, washing, cleaning, etc but do not earn money	

Table 2 shows that the disease related variables of the respondents. Majority of respondents had diagnosed with Angina Pectoris (47.0%) with duration of treatment as 1 year or above (56.0%). And majority of respondents were having continuous medical treatment (51.0%), presence of comorbidities: hypertension (71.7%), hyperlipidemia (71.3%), Diabetes Mellitus (56.7%) and respondents who did not get chance to participate in CHD awareness programs (56.0%) regarding cardiac rehabilitation.

TABLE 2 - Disease related Variables of the Respondents

n=100	
Variables	Frequency/Percent
<b>Clinical Diagnosis</b>	
Myocardial Infarction	28.0
Angina Pectoris	47.0
Ischemic heart failure	25.0
<b>Duration of Treatment</b>	
<1 year	44.0
$\geq$ 1 Years	56.0
<b>Mode of Treatment*</b>	
CMT	51.0
PI	12.0
CABG	8.0
CMT+PI	13.0
CMT+CABG	8.0
CMT+PI+CABG	8.0
<b>Presence of Comorbidities</b>	
Hypertension	71.7
Hyperlipidemia	71.3
Diabetes Mellitus	56.7
<b>Participated in CR program for CHD patient</b>	
Yes	44.0
No	56.0
*CMT=Continuous Medical Treatment, PI= Percutaneous Intervention and CABG= Coronary Artery Bypass Graft	

Table 3 shows awareness scores regarding CR, highest score of the respondents found in awareness regarding CHD (75.0%) and lowest in time and duration need for exercise per week (14.0%).

TABLE 3 - Awareness Scores Regarding Cardiac Rehabilitation of the Respondents

n=100	
Knowledge	Frequency/Percent
Awareness regarding CHD	75.0
Type of CHD	45.0
Causes of Angina	35.0
Causes of MI	44.0
Causes Heart failure	47.0
Diagnostic measures for CHD	32.0
Knowledge about stents	27.0
Need of cardiac rehabilitation for CHD patient	58.0
Type of Exercise	26.0
Time and duration need for exercise/week	14.0
Point to be considered before starting exercise	21.0
Meaning of healthy diet	63.0
Special point to maintain QOL of patients	53.0
Knowledge regarding right time for resuming duty after CABG	24.0
Precautions to be taken to avoid complication	59.0
Management of emergency conditions	49.0
Complication of CHD	53.0

Table 4 shows the respondents' level of awareness regarding CR and majority were unaware (57.0%) about CR among CHD patients.

TABLE 4 - Respondents' Level of Awareness Regarding CR

n=100	
Awareness Level	Frequency/Percent
Unaware (<57.0%)	57.0
Aware (≥57.0%)	43.0
Total	100.0
Median score of overall Awareness=17 (IQR=14.0-19.75)	

Table 5 shows the level of awareness and socio-demographic characteristics were significantly associated with age (p=0.007) and educational status (p=0.013) at p value <0.05.

TABLE 5- Association between Level of Awareness regarding CR and Socio- demographic Characteristics

n=100				
Socio-demographic characteristic	Level of Awareness regarding CR		α <sup>2</sup> Value	P Value*
	Aware No. (%)	Unaware No. (%)		
Age in years				

<54	29(55.8)	23(44.2)	7.20	0.007
54 and above	14(29.2)	34 (70.8)		
<b>Gender</b>				
Male	24 (43.6)	31 (56.4)	0.02	0.887
Female	19 (42.2)	26 (57.8)		
<b>Residence</b>				
Urban	21 (40.4)	31 (59.6)	0.302	0.582
Rural	22 (45.8)	26 (54.2)		
<b>Living Status</b>				
Living Single	8(33.3)	16(66.7)		
Living with Family	35 (46.1)	41 (53.9)	1.20	0.273
<b>*Ethnicity</b>				
Brahmin/Chhetri	26(44.1)	33(55.9)	0.067	0.796
Janajati/Dalit	17(41.5)	24(58.5)		
<b>Religion</b>				
Hindu	36 (42.9)	48(57.1)	0.004	0.947
Non-Hindu	7(43.8)	9(56.2)		
<b>Type of Family</b>				
Nuclear	20(40.8)	29 (59.2)		
Joint Family	23 (45.1)	28 (54.9)	0.187	0.665
<b>Educational status</b>				
Literate	31 (53.4)	27 (46.6)	6.151	0.013
Illiterate	12 (28.6)	30 (71.4)		
<b>Occupation</b>				
Farming & Housework	24 (42.9)	32 (57.1)	0.001	0.974
Service & Business	19(43.2)	25(56.8)		
Significance level<0.05, *Pearson chi square				

And Table 6 also represented that level of awareness and disease related variables was highly significant with duration of treatment (p=0.039) and participation in awareness program (p=0.001) at p value <0.05.

TABLE 6 - Association between Level of Awareness regarding CR and Disease Related Variables

n=100				
Disease related variables	Level of Awareness regarding CR		α <sup>2</sup> Value	P Value*
	Aware No. (%)	Unaware No. (%)		
<b>Clinical Diagnosis</b>				
Angina & IHD	33(45.8)	39(54.2)	0.842	0.359
Myocardial Infarction	10(35.7)	18(64.3)		
<b>Duration of Treatment</b>				
<1 year	24(54.5)	20(45.5)	4.273	0.039
≥1 years	19(33.9)	37(66.1)		
<b>Mode of Treatment</b>				
Invasive	21(42.9)	28(57.1)	0.001	0.977

Non- invasive	22(43.1)	29(56.9)		
Regular intake of hypertensive medication				
Yes	35(79.5)	9(20.5)	1.638	0.201
No	15(65.2)	8(34.8)		
Co-morbidity				
Present	36(42.9)	48(57.1)	0.004	0.947
Absent	7(43.8)	9(56.2)		
Participated in CR program				
Yes	10(22.7)	34(77.3)	13.175	0.001
No	33(58.9)	23(41.1)		
Significance level <0.05, *Pearson chi-square				

## Discussion

Findings of this study suggest that the mean age of respondents was 53.23 years in this study. While different studies reported slightly higher mean age such as in Canada<sup>9</sup> (64.6) and in China<sup>10</sup> (62.51), which might be due to the setting of study. In this study, regarding socio-demographic characteristics, majority of the respondents were male (55.0%), resided in urban (52.0%), living with family (76.0%), Hindu (84.0%), belonged to joint family (51.0%), and literate (58.0%). This finding was consistent with a study findings reported that the seventy two percentages of the respondents were male, residing in urban area (53.1%), living with family (92.5%), Hindu by religion (85.0%), and literate (63.4%).<sup>11</sup>

Regarding disease related variables, majority of respondents had Angina Pectoris (47.0%) with duration of treatment as 1 year or above (56.0%), continuous medical management as a mode of treatment (51.0%), presence of highest co morbidities were hypertension (71.7%) and not participating in CR program (56.0%). This findings was consistent with a study done in Nepal.<sup>12</sup> Similarly, another study conducted in Spain reported angina pectoris is the most common diagnosis among CHD patient.<sup>13</sup> Fifty seven percent of the respondents were unaware about CR. This finding was consistent with a study done in China reported by Zhou (2017) that level of awareness was low (52.69%).<sup>8</sup>

Similarly, the significant associated factors between level of awareness regarding CR and selected variables were age ( $p=0.007$ ), educational status ( $p=0.013$ ), duration of treatment of more than or equal to one year ( $p=0.039$ ) and less participation in awareness programs ( $p=0.001$ ). This finding was supported by a study done in China which found that older CHD patients were significantly associated with low level of awareness with CR program.<sup>8</sup>

## Conclusion

Our study found that patient's awareness on CR was low, especially for fifty four and above aged group, illiterate CHD patients, those receiving treatment for more than and equal to one years of duration of treatment and those that didn't participate in awareness program. Hence, health care professionals including nurses should organize & promote CR program and strengthen the counseling program to enhance their awareness level as well as for improving the quality of life.

## Limitations

It is hospital based study and sampling might also have affected the result of this study which limits the generalizability

of the findings and it does not truly represent all CHD population of Nepal.

## References

- Kayaniyiii S, Ardern C, Winstanley J, et al. Degree and correlates of cardiac knowledge and awareness among cardiac inpatients. *Patient Educ Couns* 2009; 75(1):99-107. Doi: 10.1016/j.pec.2008.09.005. <https://doi.org/10.1016/j.pec.2008.09.005>
- Nekouei ZK, Yousefy A, Doost HTN, et al. Structural Model of psychological risk and protective factors affecting on quality of life in patients with coronary heart disease: A psychocardiology model. *J Res Med Sci* 2014;19(2):90-8.
- Stafford L, Berk M, Reddy P, et al. Comorbid depression and health-related quality of life in patients with coronary artery disease. *J Psychosom Res*.2007;62(4):401-10. <https://doi.org/10.1016/j.jpsychores.2006.12.009>
- Hofer S, Saleem A, Stone J, et al. The MacNew Heart Disease Health- Related Quality of Life Questionnaire in patients with angina and patients with ischemic heart failure. *Value Health*. 2012 Jan;15(1):143-50. <https://doi.org/10.1016/j.jval.2011.07.003>
- Acharya P, Adhikari RR, Bhattarai J, et al. Delayed presentation of acute coronary syndrome:A challenge in it's earlymanagment. *J Nepal Med Assoc* 2009;48(173):1-4. <https://doi.org/10.31729/jnma.173>
- Vaidya A, Pokharel PK, Nagesh S, et al. Prevalence of coronary heart disease in the urban adult males of eastern Nepal: a population-based analytical cross-sectional study. *Indian Heart J* 2009;61(4):341-7.
- Schlitt A, Wischmann P, Wienke A, et al. Rehabilitation in Patients with Coronary Heart Disease- participation and its effect on prognosis. *Dtsch Arztebl Int* 2015; 112: 527-34. <https://doi.org/10.3238/arztebl.2015.0527>
- Zhou Y, Li J, Du S, et al. Cardiac rehabilitation knowledge in patients with Coronary heart disease in Baoding city of China: A cross- sectional study. *International Journal of Nursing Sciences* 2017 (4):24-28. <https://doi.org/10.1016/j.ijnss.2016.12.011>
- Norris CM, Ghali WA, Gallbraith PD, et al. Knudtson ML. Women with coronary artery disease report worse health-related quality of life outcomes compared to men. *Health Qual Life Outcomes*. 2004;2:1-11. <https://doi.org/10.1186/1477-7525-2-1>
- Shrestha R, Shrestha S, Adhikari B, et al. Health-Related Quality of Life and Behavioural Risk Factors among Coronary Heart Disease Patients in a Tertiary Hospital. *Int J Health Sci Res*. 2015; 5(11):211-217.
- Shrestha R & Shrestha S. Health-Related Quality of Life of Patients with Coronary Heart Disease Attending a Cardiac Care Centre in Kathmandu. *Journal of Institute f Medicine*. 2015; 37(1):4-10. <https://doi.org/10.1016/j.jacc.2015.06.529>
- Shrestha R. & Shrestha S. Effectiveness of Educational Intervention on Life Style Management among Coronary Heart Disease Patinets Attending a Cardiac Care Centre in Kathmandu. *Journal of Institute of Medicine* 2016; 38(1):20-25.
- Failde II, Soto MM. Changes in health related quality of life 3 months after an acute coronary syndrome. *BMC Public health*. 2006;6(18):1-10. <https://doi.org/10.1186/1471-2458-6-18>