

Isolated Myo-cysticercosis of the Sternocleidomastoid Muscle in a Strict Vegetarian Child: A Rare and Unusual Case Report

Sedhai M¹, Sapkota A²

¹Department of Otolaryngology, Manipal College of Medical Sciences, Pokhara, Nepal

²Department of Ophthalmology and Vitreo-retinal Surgery, B.P Koirala Lions Center for Ophthalmic Studies, Institute of Medicine, Kathmandu, Nepal

Received: March 01, 2026

Accepted: April 05, 2026

Published: August 12, 2026

Cite this paper: Sedhai M, Sapkota A. Isolated Myo-cysticercosis of the Sternocleidomastoid Muscle in a Strict Vegetarian Child: A Rare and Unusual Case Report. *Nepal Journal of Medical Sciences*. 2026;11(2):102-109. <https://doi.org/10.3126/njms.v11i2.98521>

ABSTRACT


Introduction: Cysticercosis, a parasitic disease endemic to developing countries, is caused by the larval stage of *Taenia solium*. While neurocysticercosis is common, isolated myocysticercosis, especially of the sternocleidomastoid (SCM) muscle, is exceedingly rare, particularly in pediatric patients.

Case presentation: We report the case of a 9-year-old strictly vegetarian female child presenting with painless left-sided neck swelling. Ultrasonography revealed a well-defined cyst with a central echogenic focus, suggestive of a scolex. Fine-needle aspiration cytology (FNAC) confirmed the diagnosis of cysticercosis. Computed tomography ruled out a neuro-ocular involvement. The patient was treated conservatively with albendazole, corticosteroids, and antihistamines, which led to complete resolution within three months and no residual disease or recurrence after six months.

Conclusion: This case highlights the importance of considering myocysticercosis in the differential diagnosis of neck swelling in children, even among vegetarians. Early diagnosis using noninvasive modalities and conservative management can ensure excellent outcomes while avoiding unnecessary surgical interventions in endemic regions.

Keywords: Albendazole; Case report; Cysticercosis, Myocysticercosis; Pediatric Parasitic Infection

Corresponding Author: Milan Sedhai, MS, MRCS (ENT), Lecturer, Department of Otolaryngology, Nepal Manipal College of Medical Sciences, Pokhara, Nepal. Email: milan.sedhai@nepalmanipal.edu.np



Licensed under CC BY 4.0 International License which permits use, distribution and reproduction in any medium, provided the original work is properly cited

INTRODUCTION

Cysticercosis results from tissue infection by the larval form (*Cysticercuscellulosae*) of the pork tapeworm, *Taenia solium*. The disease is most common in regions with poor sanitation and limited food safety practices, such as parts of Asia, Latin America, and sub-Saharan Africa. [1] Humans acquire infection through the ingestion of *T. solium* eggs that contaminate food or water. After ingestion, the embryos penetrate the intestinal lining, enter the bloodstream, and develop into cysticerci within the tissues.

Neurocysticercosis is the most common and severe form, but it may also lodge in the subcutaneous tissue, muscle, eyes, and heart. [1] Muscular cysticercosis is uncommon and is often asymptomatic. When symptomatic, it usually appears as a painless swelling. Sternocleidomastoid (SCM) muscle involvement is very rare, with only a few adult cases reported. [2,3,4,5,6]

Pediatric myocysticercosis is extremely rare, particularly in cases of isolated SCM without neurological symptoms. This rarity is due to a

long incubation period and underreporting. Only one pediatric case of SCM was reported by Chaudhari et al. [6] Tripathy et al. described solitary deltoid muscle cysticercosis in a child, highlighting its diagnostic challenges. [7]

High-resolution ultrasonography is the preferred diagnostic tool, showing cysts with a central scolex and surrounding inflammation. [8] Fine needle aspiration cytology (FNAC) aids in diagnosis by revealing parasitic fragments and inflammatory cells. [2, 3, 9] Treatment is mainly medical, with albendazole and corticosteroids. [1] while surgery is rarely required.

Early recognition is key to avoiding misdiagnosis and unnecessary surgery, particularly in endemic regions.

Case Report

A 9-year-old female child, strict vegetarian since birth, presented with a gradual, painless swelling in the left Level II region of the neck. There was no fever, weight loss, appetite loss, trauma, family history, drug use, contact of tuberculosis or allergies. On ENT and general examination, a single 1 × 1 cm partially mobile, fluctuant,

non-tender swelling free from overlying skin or deeper tissues, more prominent on turning of head to right was noted. Nasopharyngeal exam was unremarkable. (Fig 1A, 1B).




Fig 1: The clinical presentation of a 9-year-old female child with a single 1 × 1 cm partially mobile, fluctuant, non-tender swelling free from overlying skin or deeper tissues, more prominent on turning of head to right; A-Left lateral view, B- Oblique view (for better visualization of swelling)

Laboratory tests including complete blood count and renal function were within normal limits. Ultrasonography of the neck revealed a bulky left sternocleidomastoid muscle with a 9 × 7 mm well-defined oval cyst containing a 2 mm

echogenic focus likely representing calcification, no internal vascularity, and surrounding inflammatory changes: features suggestive of myocysticercosis of left Sternocleidomastoid region. Multiple bilateral cervical lymph nodes up to 4 × 5 mm in size showed reactive morphology. (Fig 2A)




Fig 2: (A): Ultrasonography image of the patient at the time of presentation showing- abulky left sternocleidomastoid muscle with a 9 × 7 mm well-defined oval cyst containing a 2 mm echogenic focus likely representing calcification/scolex, no internal vascularity, and surrounding inflammatory changes: features suggestive of myocysticercosis of left Sternocleidomastoid

region. Multiple bilateral cervical lymph nodes up to 4×5 mm in size showed reactive morphology;

(B): Repeat Ultrasonography image of the same patient at 1 month follow-up showing- lesion shrinkage to 8.8×4.8 mm without internal complexity (radiological signs of improvement)

A CT scan of the head and neck (axial and coronal views) was performed. IgG Serum cysticercosis antibody levels were sent. FNAC of the lesion was carried out. In addition, stool microscopy was sent that showed no ova or cysts.

FNAC cytology demonstrated parasitic structures-including scolices and hooklets-with mixed acute and chronic inflammatory cells, foamy histiocytes, eosinophils, and multinucleated giant cells in a proteinaceous background, diagnostic of cysticercosis (Fig 3). CT imaging also supported localization of the cyst within the left sternocleidomastoid muscle, with no intracranial or ocular involvement. However, IgG serum cysticercosis antibodies

were negative (value: 0.50; Normal limit: <0.90) (Fig 4). Likewise, neurology and ophthalmology evaluations were performed to exclude neuro- or ocular cysticercosis; findings were negative.

The image shows a fine needle aspiration cytology report from the Department of Lab - Cytopathology. The report includes patient information, specimen information, and a detailed microscopic description. The interpretation states: "05YRS/F, Left cervical nodule: FNAC: - Cytomorphological features are suggestive of Cysticercosis." The report is signed by a pathologist and dated 10/07/2024.

Fig 3: Fine needle aspiration cytology report of the patient confirming the diagnosis of cysticercosis

The image shows a serum IgG antibody to Cysticercosis report. The report includes patient information and a table showing the results of the test. The result is 0.50, which is below the normal limit of 0.90, indicating seronegativity. The report is signed by a pathologist and dated 10/07/2024.

Fig 4: Serum IgG antibody to Cysticercosis report of the patient showing no elevation of antibody levels (Seronegativity)

At baseline weight of 22 kg, treatment was started with albendazole 165 mg orally twice daily (2.1 ml BD) (dosage: 15 mg/kg/day) for one month, prednisolone 5 mg once daily for 15 days, and fexofenadine 120 mg at bedtime for 15 days.

At 1-month follow-up, the swelling had reduced substantially ($<0.5 \times 0.5$ cm) (Fig 5A, 5B). Ultrasonography showed lesion shrinkage to 8.8×4.8 mm without internal complexity (Fig 2B). Albendazole was continued for another one month; steroids were discontinued.




Fig 5: The clinical status of the patient after one month of medical therapy with oral albendazole and 15 days therapy with corticosteroid. The swelling has decreased to be <0.5 cm $\times$ 0.5 cm; (A): Lateral view, (B): Oblique view (for better visualization)

After two months of therapy, the swelling was clinically imperceptible. Repeat ultrasonography confirmed absence of cyst in the sternocleidomastoid muscle, with residual non-significant lymph nodes. Eventually, Albendazole was given for an additional month and then stopped. Final ultrasound at three months showed no evidence of cyst or lymphadenopathy. The patient was followed for six months post-therapy, remaining asymptomatic with no signs of recurrence (Fig 6).



Fig 6: The clinical photograph of the patient at final 6 months post therapy; there were no signs of residual pathology or recurrence

DISCUSSION

Cysticercosis remains a notable parasitic disease in areas with poor sanitation and hygiene, including Nepal, India, parts of Latin America, and sub-Saharan Africa. [1] While neurocysticercosis is the predominant presentation, muscular forms are far less common and often go unrecognizable. Within skeletal muscles, SCM involvement is rare. [2,3,4,5,6,9] and pediatric cases without neurological or ocular lesions are exceptionally uncommon; the only previous pediatric SCM case was reported by Chaudhari et al. [6]

Our case involved a 9-year-old strictly vegetarian female with a solitary, painless, gradually enlarging swelling in the left Level II cervical region. The absence of systemic features and slow progression are typical features of isolated myocysticercosis. [3,5] Her vegetarian status suggests that cysticercosis can occur via ingestion of *T. solium* eggs in contaminated food or water. [1]

High-resolution ultrasonography revealed a well-defined cyst with an echogenic nidus suggestive of a scolex or calcification, which is a

hallmark feature of cysticercosis. [8] Similar ultrasonographic patterns were noted by Thapa et al., who detected SCM involvement in an adult female. [1]

FNAC confirmed cysticercosis by identifying hooklets, scolices, and multinucleated giant cells, which is consistent with the findings of Sharma et al. and Chaudhari et al. [2,6] As noted by Marasini et al. solitary intramuscular cysticercosis can mimic a pseudotumor, making FNAC critical for avoiding unnecessary excision. [10]

We excluded systemic dissemination through CT imaging of the brain and orbits and full neurological and ophthalmological evaluations that showed no other foci. As Goyal et al. emphasized, thorough evaluation is essential, even in localized muscular cases, to rule out neurocysticercosis or ocular involvement. [3]

Due to the paucity of pediatric-specific guidelines for isolated myocysticercosis, treatment generally follows neurocysticercosis protocols. [8] Albendazole (15 mg/kg/day) for at least 14 days is preferred for the treatment of single cysts. Corticosteroids such as

prednisolone are co-administered to counteract inflammatory reactions. [1] In our patient, albendazole, prednisolone, and fexofenadine produced excellent clinical and radiological improvement, aligning with other reports. [4,5,6] At six months, the lesion completely resolved without recurrence, similar to the outcomes in adult cases and the pediatric report by Chaudhari et al. [6] Structured follow-up with serial ultrasonography ensured close monitoring and early relapse detection.

This case highlights the critical role of public health measures like improving sanitation, ensuring clean water, and promoting hygiene to prevent these parasitic infections. Educating communities on properly washing fruits and vegetables and avoiding untreated wastewater for irrigation is essential, even for vegetarian households at risk of contamination. It also emphasizes the importance of interdisciplinary collaboration among pediatricians, radiologists, pathologists, and infectious disease specialists for accurate diagnosis and management. In resource-limited settings, high-resolution

ultrasonography and FNAC are indispensable alternatives when MRI is unavailable.

CONCLUSIONS

Documenting rare pediatric cases like isolated SCM myocysticercosis enriches clinical knowledge, helps avoid unnecessary invasive procedures, and guides standardized treatment approaches.

DATA AVAILABILITY

The datasets used during the current study are available from the corresponding author on reasonable request.

ABBREVIATIONS

ENT: Ear Nose Throat

SCM: Sternocleidomastoid muscle

CT: Computed tomography

MRI: Magnetic resonance imaging

FNAC: Fine Needle Aspiration Cytology

IgG: Immunoglobulin G

CONFLICT OF INTEREST

None

SOURCES OF FUNDING

None

REFERENCES

1. Garcia HH, Gonzalez AE, Evans CA, Gilman RH. *Taenia solium* cysticercosis. *Lancet*. 2003;362(9383):547-556. [\[DOI\]](#)
2. Sharma R, Gautam P, Kumar S, Elhence P, Bansal R, Gupta G. Isolated cysticercosis cellulosae of sternocleidomastoid muscle: a case report with review of literature. *Indian Journal of Otolaryngology and Head and Neck Surgery*. 2011;63(Suppl 1):127-130. [\[DOI\]](#)
3. Goyal A, Kalra Y, Babu M, Ingale M, Chavan P. A rare presentation of cysticercosis in the sternocleidomastoid muscle. *Cureus*. 2024;16(5):e61275. [\[DOI\]](#)
4. Singh R, Kumar S, Galagali JR, Ramakrishnan N. Solitary cysticercosis of sternocleidomastoid muscle: a rare entity. *International Journal of Otorhinolaryngology and Head and Neck Surgery*. 2016;2(3):277-279. [\[DOI\]](#)
5. Hamza N, Sureshkumar S, Francis A. Cysticercosis of sternocleidomastoid muscle presenting as neck swelling: a case report. *Case Reports in Clinical Practice*. 2023;7(4):174-178. [\[DOI\]](#)
6. Chaudhari DG, Shah NR, Chaudhari AS. A rare case of an isolated muscular cysticercosis in a 5-year-old child. *International Journal of Contemporary Pediatrics*. 2024;11(11):1678-1680. [\[DOI\]](#)
7. Tripathy SK, Sen RK, Sudes P, Dhatt S. Solitary cysticercosis of deltoid muscle in a child: diagnostic dilemma. *Journal of Orthopaedics*. 2009;6(2):e11. [\[DOI\]](#)
8. Vijayaraghavan SB. Sonographic appearances in cysticercosis. *Journal of Ultrasound in Medicine*. 2004;23(3):423-427. [\[DOI\]](#)
9. Thapa S, Lamichhane N, Joshi S. Isolated cysticercosis of sternocleidomastoid muscle: role of ultrasonography. *Case Reports in Infectious Diseases*. 2021;2021:7102416. [\[DOI\]](#)
10. Marasini RP, Thapa PB, Gautam P. Solitary intramuscular cysticercosis: a rare pseudotumour. Report of two cases. *Journal of Chitwan Medical College*. 2015;5(2):42-45. [\[DOI\]](#)